

# Vignette for package `blm`

Stephanie A. Kovalchik

## Abstract

The `blm` package provides functions for fitting flexible binomial models for cohort studies of a binary outcome and population-based case-control studies. The binomial linear model (BLM) is a strictly linear model. The linear-expit (LEX-PIT) model allows risk to be expressed as a function of linear and nonlinear effects, where nonlinear effects take the form of the inverse logit function. Estimation of the model parameters is based on constrained maximum likelihood, which ensures that the fitted model yields feasible risk estimates. In this vignette, BLM and LEXPIT model fitting is demonstrated with analyses of a simulated population-based case-control study.

## Binomial linear model (BLM)

### Model

Given the binary event  $y_i$ , the probability that  $Y_i = 1$  under a binomial linear model (BLM) is a linear function of covariates  $x_i$ ,

$$\pi_i = x_i' \beta \quad (1)$$

Each  $\beta$  of nonconstant covariates represents the risk difference associated with a unit change in the given covariate, when all other factors are fixed.

Suppose that  $\tilde{x}$  is the covariate pattern for a subject from the target population of the model whose risk we want to estimate. To be a valid risk,  $\tilde{x}'\beta \in (0,1)$ . In general, we might not be able to specify all of the possible  $\tilde{x}$  of our population. Instead, we make use of the  $x_i$  from our sample and require that all  $x_i'\beta \in (0,1)$ . Thus, the set of covariate patterns of the study sample defines the *feasible region* for  $\beta$ .

To ensure that the estimates for  $\beta$  are within the region of feasibility, constrained maximum likelihood is used. Since the system of constraints are linear in the parameters, an adaptive barrier algorithm (Lange, 2010) can be used to perform the constrained optimization as implemented by `constrOptim`. For cohort studies, the objective function is a penalized binomial log-likelihood with each  $\pi_i$  defined by Equation (1). For population-based case-control studies, the objective function is a penalized pseudo-likelihood where each control subject's contribution to the binomial likelihood is weighted by a sampling weight  $w_i$  that reflects their representativeness of the target population. By definition, each case's weight is  $w_i = 1$ .

### Model fitting

As an illustration of the model syntax we consider a model to estimate the risk of disease based on a simu-

lated population-based case control study. We begin the R session by loading the package, `blm`, and the dataset `ccdata`.

```
> library(blm)
> data(ccdata)
> names(ccdata)

[1] "female" "packyear" "strata" "y" "w"

> table(ccdata$y)

 0  1
378 378
```

The sample consists of 756 subjects and case status is indicated by the variable `y`. There are two design variables, the `strata` and inverse sampling fractions `w`, and two candidate explanatory variables, which are an indicator for female gender, `female`, and a discrete variable, `packyear`, indicating the number of pack-years smoked.

The syntax for `blm` is much like `lm`, consisting of `formula` and `data` arguments. For population-based estimates, we need to additionally include the design information on sample stratification and the sampling fractions. The following code fits a population-based linear risk model with additive effects due to female gender and packyears.

```
> fit <- blm(y ~ female + packyear, data = ccdata,
+           weight = ccdata$w,
+           strata = ccdata$strata)
> fit

y ~ female + packyear
(Intercept)      female      packyear
 0.07048229  0.01110223  0.01588852

> summary(fit)

              Est.      Std. Err    t-value    p-value
(Intercept) 0.07048229 0.014534192  4.8494124 7.524509e-07
female      0.01110223 0.020695132  0.5364659 2.958975e-01
packyear    0.01588852 0.001146059 13.8636069 2.161370e-39
```

Converged: TRUE

The method `summary` provides measures of variance and Wald tests of significance for each fitted parameter. Also, a logical object indicates whether convergence of the optimization algorithm was achieved.

The variance-covariance is estimated from an influence-based method Deville (1999). The coefficients and variance-covariance can be extracted directly using `coef` and `vcov`.

```
> coef(fit)

(Intercept)      female      packyear
 0.07048229  0.01110223  0.01588852

> vcov(fit)

              [,1]      [,2]      [,3]
[1,] 2.112427e-04 -1.924858e-04 -4.816776e-06
[2,] -1.924858e-04 4.282885e-04 -1.343252e-06
[3,] -4.816776e-06 -1.343252e-06 1.313452e-06
```

Each regression coefficient for an explanatory variable of the BLM model provides an estimate of the adjusted risk difference associated with a unit increase in the given variable. Thus, the fitted model suggests that there is a 1.1% increased risk for females and a 15.9% increased absolute risk for every 10-year increase in cumulative pack-years smoked. To obtain confidence intervals for these parameters, we can use the `confint` method.

```
> confint(fit)
              Est.      Lower      Upper
(Intercept) 0.07048229 0.04199580 0.09896878
female       0.01110223 -0.02945948 0.05166395
packyear     0.01588852 0.01364228 0.01813475

> confint(fit, parm="female")
              Est.      Lower      Upper
[1,] 0.01110223 -0.02945948 0.05166395
```

To assess the constraints imposed on the model, we can examine the `barrier.value`, which is one of the slots of the `blm` class.

```
> fit@barrier.value
[1] 0.04078781
```

Risk estimates near the boundary could be an indication of influential points or a poor-fitting model with BLM or LEXPIT. Boundary estimates for a given distance criterion can be obtained with the function `which.at.boundary`.

```
> which.at.boundary(fit)
No boundary constraints using the given criterion.

> which.at.boundary(fit, criter = 1e-3)
No boundary constraints using the given criterion.
```

In the above, we first use a default criterion of  $1e-06$  or  $0.999999$ . In the second case, we provide a user-specified criterion. With either criterion, no estimate was at the boundary.

As an additional way to evaluate potential influential points in a BLM or LEXPIT model fit, each class has a method `dBeta`, which returns the jackknife deviates by subject and parameter. Here, we plot a sparkline-like figure of the `packyear` deviates by subject (Figure 1).

```
> plot(y = dBeta(fit)[,2], x = 1:nrow(ccdata),
+      type = "h")
```




Figure 1: Jackknife deviates for variable `packyear` by subject

There are several functions to evaluate the BLM model fit. McFadden's R-squared, adjusted and unadjusted, provides a measure of the variability explained by the explanatory variables.

```
> Rsquared(fit)
$R2
[1] 0.1643529
$R2adj
[1] 0.1606472
```

Comparisons of observed and expected counts for the target population can be made with the function `EO` for expected and observed. A factor can also be supplied to compare the expected to observed within subgroups defined by the categorical variable.

```
> EO(fit)
              E  0      Eto0  lowerCI  upperCI
Overall 387.2257 378 1.024407 0.9261713 1.133062
```

```
> EO(fit, ccdata$female)
              E  0      Eto0  lowerCI  upperCI
0 192.5145 183 1.051992 0.9101015 1.216004
1 194.7112 195 0.998519 0.8677618 1.148979
```

When the number of covariate classes represented by the model are few, we can perform a goodness-of-fit test with Pearson's chi-squared statistic. This compares the observed to expected within each unique risk type defined by the model.

```
> gof.pearson(fit)
              E  0
1 17.22 22
2 23.27 16
3 79.42 59
4 72.60 86
5 23.25 24
6 18.30 13
7 83.50 78
8 69.65 80
```

```
Chi-squared: 42.24285
P-value: 1.646512e-07
```

This suggest a lack of fit at the population level. When a model has more than 10 covariate classes, the Hosmer-Lemeshow test should be used. For the `blm` function, this test is implemented by the function `gof`.

## Linear-Expit (LEXPIT) model

### Model

Suppose we were concerned that the linear assumption for pack-years might not be valid. If we thought that linearity on the relative risk scale was a more plausible model for the effect of pack-years, we could consider a LEXPIT model. The LEXPIT model describes the probability of  $Y_i = 1$  as a function of linear and nonlinear effects, where the nonlinear effects are the expit function (the inverse of the logit),  $\text{expit}(x) = \exp(x)/(1 + \exp(x))$ .

$$\pi_i = x_i'\beta + \text{expit}(z_i'\gamma) \tag{2}$$

The  $x_i$  variables are linear effects and  $z_i$  are the logistic effects. The first component of  $z_i$  is an intercept term, so that when the remaining components are 0,  $\text{expit}(\gamma_0)$  is the baseline risk. As in BLM,  $\beta$  represent risk differences for unit changes in  $x_i$ . The coefficients  $\gamma$  are odds ratios after baseline adjustment for the effects of  $x_i'\beta$ , what we can think of as ‘residual odds ratios’.

The LEXPIT model provides a more flexible way to estimate risk differences since it imposes fewer parameter constraints. This is possible because any  $z_i'\gamma$  yields a probability measure.

Estimation for the LEXPIT model proceeds in two stages. The first stage fixes the expit parameters and estimates the linear coefficients with constrained maximization as described for the BLM in Section 1. Thus, in this stage, the expit term can be thought of as an offset in a BLM model. The second stage maximizes the expit parameters treating the linear term as fixed. Maximization at this stage uses a standard iterative reweighted least squares algorithm with modified weights that incorporate the linear risk offset.

### Model fitting

The syntax for the `lexpit` takes two formula arguments: one for the linear components and one for the expit components. Note that the intercept is always included in the expit term. Otherwise, the syntax is identical to `blm`.

```
> fit.lexpit <- lexpit(y~female, y~packyear,
+                   data = ccddata,
+                   weight = ccddata$w,
+                   strata = ccddata$strata)
> summary(fit.lexpit)
```

```
Linear effects:
      Est.   Std. Err t-value  p-value
female 0.01922563 0.01540602 1.24793 0.1062224
```

```
Expit effects:
      Est.   Std. Err  t-value    p-value
(Intercept) -2.6394437 0.169602864 -15.56249 7.395456e-48
packyear     0.1015473 0.007184217  14.13477 1.047133e-40
```

```
Converged: TRUE
```

All of the methods described for the `blm` class are also available for `lexpit` objects.

```
> which.at.boundary(fit.lexpit)
```

No boundary constraints using the given criterion.

```
> confint(fit.lexpit)
```

	Est.	Lower	Upper
female	0.01922563	-0.01096962	0.04942088
(Intercept)	-2.63944371	-2.97185921	-2.30702821
packyear	0.10154727	0.08746646	0.11562807

```
> gof.pearson(fit.lexpit)
```

	E	O
1	16.28	22
2	16.70	16
3	72.09	59
4	79.67	86
5	24.47	24
6	13.99	13
7	77.70	78
8	77.31	80

```
Chi-squared: 11.99811
```

```
P-value: 0.06201106
```

The goodness-of-fit has improved with the LEXPIT model, suggesting that multiplicative rather than linear risk effects might be a more suitable model for the effect of continuous pack-years.

## Conclusion

The `blm` package provides two models, BLM and LEXPIT, that can be used to obtain direct estimates of absolute risk and risk differences for binary data obtained from observational study designs. Fitting the `blm` and `lexpit` models will be straight-forward for R users because they provide similar syntax and methods as the `lm` class. The BLM and LEXPIT models provide alternatives to logistic regression analysis of binary data that are appealing for epidemiological interpretation because they allow for the assessment of risk associations on an absolute rather than relative risk scale.

## Bibliography

Deville JC (1999). Variance estimation for complex statistics and estimators: linearization and residual techniques *Survey methodology* 25(2):193–204

Lange K (2010). *Numerical Analysis for Statisticians*. Springer-Verlag, New York.

Madsen K, Nielsen H, and Tingleff O (2004). *Optimization with constraints*. IMM, Technical University of Denmark.