

## APPROXIMATING DISTRIBUTION FUNCTIONS BY ITERATED FUNCTION SYSTEMS\*

STEFANO M. IACUS<sup>1</sup> AND DAVIDE LA TORRE<sup>1</sup>

**Abstract.** In this small note an iterated function system on the space of distribution functions is built. The inverse problem is introduced and studied by convex optimization problems. Some applications of this method to approximation of distribution functions, to distribution function estimation and randomly censored survival analysis are given. These statistical applications are special cases in which there is no need to solve an optimization problem to obtain good approximations in terms of statistical quantities.

**Résumé.** Dans ce petit travail on construit un système de fonctions itérées dans l'espace de fonctions de répartition. Le problème inverse est introduit et étudié dans le cadre d'un problème d'optimisation. Quelques applications de cette méthode à l'approximation des fonctions de distribution, à l'estimation d'une fonction de répartition et à l'analyse de données de durée sont présentées. Ces derniers problèmes d'estimation sont des cas particuliers pour lesquels il n'y a pas besoin de résoudre un problème d'optimisation pour obtenir de bonnes approximations par rapport à des mesures statistiques.

**1991 Mathematics Subject Classification.** 62E17, 62H10, 37H.

## 1. INTRODUCTION

The Iterated Function Systems (IFSs) were born in mid eighties [2, 7] as applications of the theory of discrete dynamical systems and as useful tools to build fractals and other similar sets. Some possible applications of IFSs can be found in image processing theory [6], in the theory of stochastic growth models [13], and in the theory of random dynamical systems [1, 4, 9]. The fundamental result [2] on which the IFS method is based is Banach theorem.

In practical applications a crucial problem is the so-called *inverse problem*. This can be formulated as follows: given  $f$  in some metric space  $(S, d)$ , find a contraction  $T : S \rightarrow S$  that admits a unique fixed point  $\tilde{f} \in S$  such that  $d(f, \tilde{f})$  is small enough. In fact if one is able to solve the inverse problem with arbitrary precision, it is possible to identify  $f$  with the operator  $T$  which has it as fixed point.

The paper is organized as follows: Section 2 is devoted to introduce a contractive operator  $T$  on the space of distribution functions while, in Section 3, the inverse problem for  $T$  is studied in details. Section 4 is divided

---

*Keywords and phrases:* iterated function systems, optimization, distribution function estimation

\* CNR grant n.

<sup>1</sup> Dipartimento di Economia e Politica Aziendale, Via Conservatorio 7, I-20122 Milano, Italy; e-mail: [stefano.iacus@unimi.it](mailto:stefano.iacus@unimi.it); e-mail: [davide.latorre@unimi.it](mailto:davide.latorre@unimi.it).

into two parts: in the first some examples of inverse problems are analyzed and explicit solutions are given. In the second one, we introduce an estimator of the unknown distribution function based on IFSs.

## 2. A CONTRACTION ON THE SPACE OF DISTRIBUTION FUNCTIONS

Let us denote by the space of distribution functions  $F$  on  $[0, 1]$  by  $\mathcal{F}([0, 1])$  and by  $\mathcal{B}([0, 1])$  the space of real bounded functions on  $[0, 1]$ . Let us further define, for  $F, G \in \mathcal{B}([0, 1])$ ,  $d_\infty(F, G) = \sup_{x \in [0, 1]} |F(x) - G(x)|$ . So that  $(\mathcal{F}([0, 1]), d_\infty)$  is a complete metric space.

Let  $N \in \mathbb{N}$  be fixed and let:

- i)  $w_i : [a_i, b_i] \rightarrow [c_i, d_i] = w_i([a_i, b_i])$ ,  $i = 1, \dots, N-1$ ,  $w_N : [a_N, b_N] \rightarrow [c_N, d_N]$ , with  $a_1 = c_1 = 0$  and  $b_N = d_N = 1$ ;
- ii)  $w_i$ ,  $i = 1 \dots N$ , are increasing and continuous;
- iii)  $\bigcup_{i=1}^{N-1} [c_i, d_i] \cup [c_N, d_N] = [0, 1]$ ;
- iv) if  $i \neq j$  then  $[c_i, d_i) \cap [c_j, d_j) = \emptyset$ .
- iv)  $p_i \geq 0$ ,  $i = 1, \dots, N$ ,  $\delta_i \geq 0$ ,  $i = 1 \dots N-1$ ,  $\sum_{i=1}^N p_i + \sum_{i=1}^{N-1} \delta_i = 1$ .

On  $(\mathcal{F}([0, 1], d_\infty))$  we define an operator in the following way:

$$TF(x) = \begin{cases} p_1 F(w_1^{-1}(x)), & x \in [c_1, d_1) \\ p_i F(w_i^{-1}(x)) + \sum_{j=1}^{i-1} p_j + \sum_{j=1}^{i-1} \delta_j, & x \in [c_i, d_i), i = 2, \dots, N-1 \\ p_N F(w_N^{-1}(x)) + \sum_{j=1}^{N-1} p_j + \sum_{j=1}^{N-1} \delta_j, & x \in [c_N, d_N] \end{cases} \quad (1)$$

where  $F \in \mathcal{F}([0, 1])$ . In many practical cases  $w_i$  are affine maps. The new distribution function  $TF$  is union of distorted copies of  $F$ ; this is the fractal nature of the operator.

A similar approach has been discussed in [10] but here a more general operator is defined.

We stress here that in the following we will think to the maps  $w_i$  and to the parameters  $\delta_j$  as fixed while the parameters  $p_i$  have to be chosen. To put in evidence the dependence of the operator  $T$  on the vector  $p = (p_1, \dots, p_N)$  we will write  $T_p$  instead of  $T$ .

In Remark 2.2 the hypotheses ii) and iv) will be weakened to allow more general functionals.

**Theorem 2.1.**  $T_p$  is an operator from  $\mathcal{F}([0, 1])$  to itself.

*Proof.* It is trivial that  $T_p F(0) = 0$  and  $T_p F(1) = 1$ . Furthermore if  $x_1 > x_2$ , without loss of generality, we will consider the two cases:

- : i)  $x_1, x_2 \in w_i([a_i, b_i])$ ;
- : ii)  $x_1 \in w_{i+1}([a_{i+1}, b_{i+1}])$  and  $x_2 \in w_i([a_i, b_i])$ .

In case i), recalling that  $w_i$  are increasing maps, we have:

$$\begin{aligned} T_p F(x_1) &= p_i F(w_i^{-1}(x_1)) + \sum_{j=1}^{i-1} p_j + \sum_{j=1}^{i-1} \delta_j \\ &\geq p_i F(w_i^{-1}(x_2)) + \sum_{j=1}^{i-1} p_j + \sum_{j=1}^{i-1} \delta_j \\ &= T_p F(x_2) \end{aligned}$$

In case *ii*) we obtain:

$$\begin{aligned} T_p F(x_1) - T_p F(x_2) &= p_i + \delta_{i-1} + p_{i+1} F(w_{i+1}^{-1}(x_1)) - p_i F(w_i^{-1}(x_2)) \\ &= p_i(1 - F(w_i^{-1}(x_2))) + p_{i+1} F(w_{i+1}^{-1}(x_1)) + \delta_{i-1} \geq 0 \end{aligned}$$

since  $p_i \geq 0$ ,  $\delta_i \geq 0$  and  $0 \leq F(y) \leq 1$ . Finally, one can prove without difficulties the right continuity of  $T_p f$ .  $\square$

The following remark will be useful for the applications in Section 4.

**Remark 2.2.** If hypotheses i) and iv) are replaced by the following

- i')  $w_i(x) = x$ ,  $a_i = c_i$ ,  $b_i = d_i$ ,  $i = 1, \dots, N$ ,
- iv')  $p_i = p$ ,  $\delta_i \geq -p$ ,  $i = 1, \dots, N$ ,  $Np + \sum_{i=1}^{N-1} \delta_i = 1$ ,

then  $T_p : \mathcal{F}([0, 1]) \rightarrow \mathcal{F}([0, 1])$ .

**Theorem 2.3.** *If  $c = \max_{i=1, \dots, N} p_i < 1$ , then  $T_p$  is a contraction on  $(\mathcal{F}([0, 1]), d_\infty)$  with contractivity constant  $c$ .*

*Proof.* Let  $F, G \in (\mathcal{F}([0, 1]), d_\infty)$  and let it be  $x \in w_i([a_i, b_i])$ . We have

$$|T_p F(x) - T_p G(x)| = p_i |F(w_i^{-1}(x)) - G(w_i^{-1}(x))| \leq c d_\infty(F, G).$$

This implies  $d_\infty(T_p F, T_p G) \leq c d_\infty(F, G)$ .  $\square$

The following theorem states that small perturbations of the parameters  $p_i$  produce small variations on the fixed point of the operator.

**Theorem 2.4.** *Let  $\mathbf{p}, \mathbf{p}^* \in \mathbb{R}^N$  such that  $T_p F_1 = F_1$  and  $T_{p^*} F_2 = F_2$ . Then*

$$d_\infty(F_1, F_2) \leq \frac{1}{1-c} \sum_{j=1}^N |p_j - p_j^*|$$

where  $c$  is the contractivity constant of  $T_p$ .

*Proof.* In fact, recalling that  $w_i$  and  $\delta_i$  are fixed, we have

$$\begin{aligned} d_\infty(F_1, F_2) &= d_\infty(T_p F_1, T_p F_2) \\ &= \max_{i=1, \dots, N} \sup_{x \in [c_i, d_i]} \left\{ \left| p_i F_1(w_i^{-1}(x)) + \sum_{j=1}^{i-1} p_j - p_i^* F_2(w_i^{-1}(x)) + \sum_{j=1}^{i-1} p_j^* \right| \right\} \\ &\leq \sum_{i=1}^N |p_i - p_i^*| + c d_\infty(F_1, F_2), \end{aligned}$$

since

$$\begin{aligned}
& \left| p_i F_1(w_i^{-1}(x)) + \sum_{j=1}^{i-1} p_j - p_i^* F_2(w_i^{-1}(x)) + \sum_{j=1}^{i-1} p_j^* \right| \\
& \leq \sum_{j=1}^{i-1} |p_j - p_j^*| + |p_i F_1(w_i^{-1}(x)) - p_i F_2(w_i^{-1}(x))| + |p_i F_2(w_i^{-1}(x)) - p_i^* F_2(w_i^{-1}(x))| \\
& \leq \sum_{j=1}^{i-1} |p_j - p_j^*| + p_i d_\infty(F_1, F_2) + |p_i - p_i^*| \\
& \leq c d_\infty(F_1, F_2) + \sum_{j=1}^N |p_j - p_j^*|.
\end{aligned}$$

□

### 3. THE INVERSE PROBLEM AS A CONVEX CONSTRAINED OPTIMIZATION PROBLEM

Choose  $F \in (\mathcal{F}([0, 1]), d_\infty)$ . The aim of solving the inverse problem is to find a contractive map  $T : \mathcal{F}([0, 1]) \rightarrow \mathcal{F}([0, 1])$  which has a fixed point “near” to  $F$ . In fact if it is possible to solve the inverse problem with an arbitrary precision one can identify the operator  $T$  with its fixed point. With a fixed system of maps  $w_i$  and parameters  $\delta_j$ , the inverse problem can be solved, if it is possible, by using the parameters  $p_i$ . These have to be chosen in the following convex set:

$$C = \left\{ \mathbf{p} \in \mathbb{R}^N : p_i \geq 0, i = 1, \dots, N, \sum_{i=1}^N p_i = 1 - \sum_{i=1}^{N-1} \delta_i \right\},$$

We have the following result.

**Theorem 3.1.** *Choose  $\epsilon > 0$  and  $\mathbf{p} \in C$  such that  $p_i \cdot p_j > 0$  for some  $i \neq j$ . If  $d_\infty(T_p F, F) \leq \epsilon$ , then:*

$$d_\infty(F, \tilde{F}) \leq \frac{\epsilon}{1 - c},$$

where  $\tilde{F}$  is the fixed point of  $T_p$  on  $\mathcal{F}([0, 1])$  and  $c = \max_{i=1, \dots, N} p_i$  is the contractivity constant of  $T_p$ .

*Proof.* The assumptions imply  $c < 1$ . So we have:

$$d_\infty(F, \tilde{F}) \leq d_\infty(F, T_p F) + d_\infty(T_p F, T_p \tilde{F}) \leq \epsilon + c d_\infty(F, \tilde{F})$$

and so we get the assertion. □

If we wish to find an approximated solution of the inverse problem, we have to solve the following constrained optimization problem:

$$(\mathbf{P}) \quad \min_{\mathbf{p} \in C} d_\infty(T_p F, F)$$

It is clear that the ideal solution of  $(\mathbf{P})$  consists of finding a  $\mathbf{p}^* \in C$  such that  $d_\infty(T_{\mathbf{p}^*} F, F) = 0$ . In fact this means that, given a distribution function  $F$ , we have found a contractive map  $T_p$  which has exactly  $F$  as fixed point. Indeed the use of Theorem 3.1 gives us only an approximation of  $F$ . This can be improved, once fixed the maps  $w_i$ , by increasing the number of parameters  $p_i$ .

The following result proves the convexity of the function  $D(\mathbf{p}) = d_\infty(T_p F, F)$ ,  $\mathbf{p} \in \mathbb{R}^N$ .

**Theorem 3.2.** *The function  $D(\mathbf{p}) : \mathbb{R}^N \rightarrow \mathbb{R}$  is convex.*

*Proof.* If we choose  $\mathbf{p}_1, \mathbf{p}_2 \in \mathbb{R}^N$  and  $\lambda \in [0, 1]$  then:

$$D(\lambda \mathbf{p}_1 + (1 - \lambda) \mathbf{p}_2) = \sup_{x \in [0,1]} |T_{\lambda p_1 + (1-\lambda)p_2} F(x) - F(x)| \leq$$

$$\lambda \sup_{x \in [0,1]} |T_{p_1} F(x) - F(x)| + (1 - \lambda) \sup_{x \in [0,1]} |T_{p_2} F(x) - F(x)| = \lambda D(p_1) + (1 - \lambda) D(p_2).$$

□

Hence for solving problem (P) one can recall classical results about convex programming problems (see for instance [14]). A necessary and sufficient condition for  $\mathbf{p}^* \in C$  to be a solution of (P) can be given by Kuhn-Tucker conditions.

#### 4. INVERSE PROBLEM FOR DISTRIBUTION FUNCTIONS AND APPLICATIONS

In this section we consider different problems. We show that for a particular class of distribution functions the inverse problem can be solved exactly without solving any optimization problem. Then we discuss two ways of construct IFS to approximate a distribution function  $F$  with a finite number of parameters  $p_i$  and maps  $w_i$ .

As is usual in statistical applications, given a sample of  $n$  independent and identically distributed observations,  $(x_1, x_2, \dots, x_n)$ , drawn from an unknown distribution function  $F$ , one can easily construct the empirical distribution function (e.d.f.)  $\hat{F}_n$  that reads

$$\hat{F}_n(x) = \frac{1}{n} \sum_{i=1}^n \chi_{(-\infty, x]}(x_i), \quad x \in \mathbb{R},$$

where  $\chi_A$  is the indicator function of the set  $A$ . Asymptotic properties of optimality of  $\hat{F}_n$  as estimator of the unknown  $F$  when  $n$  goes to infinity are well known and studied [11, 12]. This function has an IFS representation that is exact and can be found without solving any optimization problem. We assume that the  $x_i$  in the sample are all different (this assumption is natural if  $F$  is a continuous distribution function). Let  $w_i(x) : [x_{i-1}, x_i) \rightarrow [x_{i-1}, x_i)$ , when  $i = 1 \dots n$  and  $w_1(x) : [0, x_1) \rightarrow [0, x_1)$ ,  $w_{n+1}(x) : [x_n, x_{n+1}] \rightarrow [x_n, x_{n+1}]$ , with  $x_0 = 0$  and  $x_{n+1} = 1$ . Assume also that every map is of the form  $w_i(x) = x$ . If we choose  $p_i = \frac{1}{n}$ ,  $i = 2 \dots n+1$ ,  $p_1 = 0$  and

$$\delta_1 = \frac{n-1}{n^2}, \quad \delta_i = -\frac{1}{n^2}$$

then the following representation holds:

$$T_p \hat{F}_n(x) = \begin{cases} 0, & i = 1 \\ \frac{1}{n} \hat{F}_n(x) + \frac{n-1}{n^2}, & i = 2 \\ \frac{1}{n} \hat{F}_n(x) + \frac{i-1}{n} + \frac{n-i+1}{n^2}, & i = 3, \dots, n+1. \end{cases}$$

when  $x \in [x_{i-1}, x_i)$ . Furthermore, from Corollary 2.2 we are guaranteed that

$$\lim_{s \rightarrow \infty} d_\infty(T_p^{(s)} u, \hat{F}_n) \rightarrow 0, \quad \forall u \in \mathcal{F}[0, 1].$$

Note that from the point of view of applications, constructing the e.d.f. or iterate the IFS with the given maps is exactly equivalent if one start, for example, with a uniform distribution on  $[0, 1]$  in the first iteration. So this is just a case when we can present an IFS that gives exact result for this particular class of distribution functions.

What follows, on the contrary, is more attractive from the point of view of applications. Suppose that one knows the distribution function  $F$  and wants to construct the IFS which has  $F$  as fixed point. In general one has to provide an infinite set of affine maps  $\{w_i, i \in \mathbb{N}\}$  and solve an extremal problem to find the corresponding sequence of weights  $p_i, i \in \mathbb{N}$ . This problem has not a general solution but at the same time the solution in terms of a finite, possibly few, number of maps and weights is crucial in applications like image compression and transmission.

The idea is the following: one can think at  $n$  points  $(x_1, x_2, \dots, x_n)$  as they were drawn from the distribution function  $F$ , and use the same maps  $w_i$  of the e.d.f.  $\hat{F}_n$ , then instead of using the  $p_i$  equal to  $1/n$  one solve the extremal problem as it is usual in IFS application. The corresponding IFS should have a fixed point that is a "good" approximation on  $F$ . So it is sufficient to store the simulated data and the weights instead of  $F$  itself.

We take the functional  $T_p F$  with the particular choice of  $\delta_i = 0$ . This choice is in principle not necessary but simplifies the solution of the problem. We simulated  $n$  i.i.d. observations from the distribution function  $F$  and we use the maps of the e.d.f. above.

We now try to solve the extremal problem

$$\min d_\infty(T_p F, F)$$

under the constrain  $\sum_{i=1}^n p_i = 1, p_i \geq 0, i = 1, \dots, n$  (with some  $p_i > 0$ ). The optimal solution will be  $\{\hat{p}_i, i = 1, \dots, n\}$  such that  $d_\infty(T_{\hat{p}} F, F) = 0$  that it is true in at least one case: if  $F$  equals  $\hat{F}_n$  and  $p_i = 1/n$ . Otherwise we will obtain some positive number. That means that, in principle, in the worst case we can approximate  $F$  with its empirical distribution function  $\hat{F}_n$ . But we can generally do better. So let us solve the problem: let us fix  $x_0 = 0$  and  $x_{n+1} = 1$ , then

$$\begin{aligned} d_\infty(T_p F, F) &= \sup_{x \in [0,1]} |T_p F(x) - F(x)| \\ &= \max_{i=1, \dots, n+1} \left\{ \sup_{[x_{i-1}, x_i]} |T_p F(x) - F(x)| \right\} \\ &= \max_{i=1, \dots, n+1} \left\{ \sup_{[x_{i-1}, x_i]} \left| \sum_{j=1}^{i-1} p_j - (1-p_i)F(x) \right| \right\} \\ &= \max_{i=1, \dots, n+1} \left\{ \left| \sum_{j=1}^{i-1} p_j - (1-p_i)F(x_{i-1}) \right|, \left| \sum_{j=1}^{i-1} p_j - (1-p_i)F(x_i^-) \right| \right\} \end{aligned}$$

and the last line is due to the non-decreasing property of  $F$ .

**Example 4.1.** Suppose that  $F$  is the distribution function of a uniform distribution on  $[0,1]$  and suppose that we can only draw one observation from  $F$  (or choose a point)  $x_1$ . The empirical distribution function  $\hat{F}_1(x) = \chi_{\{x_1, 1\}}(x)$  is useless if we have in mind to approximate  $F$ . Let us use the second technique: fix  $w_1 : [0, x_1] \rightarrow [0, x_1]$  and  $w_2 : [x_1, 1] \rightarrow [x_1, 1]$ . We try to solve the above extremal problem.

$$\begin{aligned} d_\infty(T_p F, F) &= \max_{i=1, \dots, n+1} \left\{ \left| \sum_{j=1}^{i-1} p_j - (1-p_i)F(x_i) \right|, \left| \sum_{j=1}^{i-1} p_j - (1-p_i)F(x_{i+1}) \right| \right\} \\ &= \max \left\{ \left| -(1-p_1) \cdot 0 \right|, \left| -(1-p_1)x_1 \right|, \left| p_1 - (1-p_2)x_1 \right|, \left| p_1 - (1-p_2) \right| \right\} \\ &= \max\{0, x_1(1-p_1), p_1(1-x_1), 0\}, \end{aligned}$$

$x_1 \in (0, 1)$ ,  $p_1 + p_2 = 1$ . Now minimizing with respect to  $p_1$  and  $p_2$  under the constraint  $p_1 + p_2 = 1$  one obtains simply  $p_1 = x_1$ . The resulting functional will be

$$T_{x_1}u(x) = \begin{cases} x_1 u(x), & x \in [0, x_1) \\ (1 - x_1)u(x) + x_1, & x \in [x_1, 1] \end{cases}$$

and it is clear that  $T_{x_1}u(x) = T_{x_1}x$ , for one iteration only, is closer than  $\hat{F}_1$  to  $F(x) = x$  and the approximation is better and better as  $x_1 \rightarrow 0$  or  $x_1 \rightarrow 1$ .

We propose now a more efficient method to approximate  $F$  when  $F$  is not to be estimated. We have already mentioned that the e.d.f. is the better estimator of an unknown distribution function  $F$ , so one can think to sample  $n$  points from  $F$  and use their values to approximate  $F$  by  $\hat{F}_n$ . As  $n \rightarrow \infty$ , the statistical literature assures almost sure convergence of  $\hat{F}_n(x)$  to  $F(x)$  for every  $x$ . We also have shown the exact IFS representation of  $\hat{F}_n$ . But this method is not efficient. On the contrary, suppose that  $F$  is a continuous distribution function. As we know  $F$ , we can think to approximate it by means of continuous functions instead of simple functions like  $\hat{F}_n$ . Choose  $n$  points  $(x_1, \dots, x_n)$  and assume that  $x_0 = 0$  and  $x_{n+1} = 1$ . One can construct the following functional

$$T_F u(x) = (F(x_i) - F(x_{i-1}))u\left(\frac{x - x_{i-1}}{x_i - x_{i-1}}\right) + F(x_{i-1}), \quad x \in [x_{i-1}, x_i),$$

$i = 1, \dots, n + 1$ . Notice that  $T_F$  is a particular case of (1) where  $p_i = F(x_i) - F(x_{i-1})$ ,  $\delta_i = 0$  and  $w_i(x) : [0, 1] \rightarrow [x_{i-1}, x_i)$ . This is a contraction and, at each iteration,  $T_F$  passes exactly through the points  $F(x_i)$ . It is almost evident that, when  $n$  increases the fixed point of the above functional will be ‘‘close’’ to  $F$ . So again, instead of sending an infinite set of weights and maps, one can send  $n$  points and the values of  $F$  evaluated at these points. All in summary, only  $2 \cdot n$  informations should be sent to reconstruct  $F$ .

For  $n$  small, the choice of a good grid of point is critical. So one question arises: how to choose them? One can proceed case by case but as  $F$  is a distribution function one can use its properties. We propose the following solution: take  $n$  points  $(u_1, u_2, \dots, u_n)$  equally spaced  $[0, 1]$  and define  $x_i = F^{-1}(u_i)$ ,  $i = 1, \dots, n$ . The points  $x_i$  are just the quantiles of  $F$ . In this way, it is assured that the profile of  $F$  is followed as smooth as possible. In fact, if two quantiles  $x_i$  and  $x_{i+1}$  are relatively distant each other, then  $F$  is slowly increasing in the interval  $(x_i, x_{i+1})$  and viceversa. This method is more efficient than simply taking equally spaced points on  $[0, 1]$ . If this method of choosing the points is used, then the functional simply reads

$$T_F u(x) = \frac{1}{n}u\left(\frac{x - x_{i-1}}{x_i - x_{i-1}}\right) + \frac{i - 1}{n}, \quad x \in [x_{i-1}, x_i), \quad i = 1, \dots, n + 1.$$

And this suggests an empirical estimator of  $F$ . If  $\hat{q}_i$ ,  $i = 1, 2, \dots, k$ ,  $k < n$ , are the empirical quantiles of the sample  $(x_1, x_2, \dots, x_n)$  of order  $\frac{i}{k}$ , then an estimator of the unknown distribution function  $F$  should be written as

$$\tilde{F}_{(k)}u(x) = \frac{1}{k}u\left(\frac{x - \hat{q}_i}{\hat{q}_{i+1} - \hat{q}_i}\right) + \frac{i - 1}{k}, \quad x \in [\hat{q}_i, \hat{q}_{i+1}),$$

$i = 1, \dots, k$ , with  $\hat{q}_0 = 0$  and  $\hat{q}_{k+1} = 1$ . As  $n$  and  $k = k(n)$  go to infinity  $\tilde{F}_{(k)}$  converges to  $F$ . Relative efficiency of  $\tilde{F}_{(k)}$  with respect to  $\hat{F}_n$  is investigated via simulations. The results are reported in Table 1 for differently shaped distribution functions and sample sizes. What emerges is that  $\tilde{F}_{(k)}$  is equivalent to the e.d.f. in the sense of the sup-norm.

#### 4.1. Applications to survival analysis

Let  $T$  denote a random lifetime (or time until failure) with distribution function  $F$ . On the basis of a sample of  $n$  independent replications of  $T$  the object of inference are usually quantities derived from the so-called survival function  $S(t) = 1 - F(t) = P(T < t)$ . If  $F$  has a density  $f$  then it is possible to define the hazard

| number of points<br>drawn from $F$ | $d_\infty \left( \tilde{F}_{(k)}^{(4)} u, F \right)$<br>(a) | $d_\infty \left( \hat{F}_n, F \right)$<br>(b) | $\frac{(a)}{(b)} \cdot 100\%$ | distribution $F$ |
|------------------------------------|---|---|-------------------------------|------------------|
| 10                                 | 0.20232   | 0.24103                                       | 83.94%                        | Beta(2,2)        |
| 50                                 | 0.09376   | 0.10241                                       | 91.56%                        | Beta(2,2)        |
| 100                                | 0.06989   | 0.07131                                       | 98.01%                        | Beta(2,2)        |
| 500                                | 0.02884   | 0.02917                                       | 98.87%                        | Beta(2,2)        |
| 1000                               | 0.02475   | 0.02506                                       | 98.78%                        | Beta(2,2)        |
| 10                                 | 0.18747   | 0.19472                                       | 96.27%                        | Beta(3,3)        |
| 50                                 | 0.09945   | 0.09777                                       | 101.72%                       | Beta(3,3)        |
| 100                                | 0.07103   | 0.07521                                       | 94.44%                        | Beta(3,3)        |
| 500                                | 0.03077   | 0.03061                                       | 100.52%                       | Beta(3,3)        |
| 1000                               | 0.01993   | 0.02018                                       | 98.74%                        | Beta(3,3)        |
| 10                                 | 0.20842   | 0.22220                                       | 93.80%                        | Beta(5,3)        |
| 50                                 | 0.10615   | 0.10517                                       | 100.93%                       | Beta(5,3)        |
| 100                                | 0.06881   | 0.07096                                       | 96.96%                        | Beta(5,3)        |
| 500                                | 0.02959   | 0.02971                                       | 99.60%                        | Beta(5,3)        |
| 1000                               | 0.02176   | 0.02194                                       | 99.17%                        | Beta(5,3)        |
| 10                                 | 0.23054   | 0.23301                                       | 98.94%                        | Beta(3,5)        |
| 50                                 | 0.08993   | 0.089347                                      | 100.66%                       | Beta(3,5)        |
| 100                                | 0.06541   | 0.06515                                       | 100.40%                       | Beta(3,5)        |
| 500                                | 0.02978   | 0.03015                                       | 98.80%                        | Beta(3,5)        |
| 1000                               | 0.01978   | 0.02003                                       | 98.77%                        | Beta(3,5)        |
| 10                                 | 0.20522   | 0.24492                                       | 83.79%                        | Beta(1,1)        |
| 50                                 | 0.10456   | 0.11990                                       | 87.20%                        | Beta(1,1)        |
| 100                                | 0.07621   | 0.08124                                       | 93.81%                        | Beta(1,1)        |
| 500                                | 0.02938   | 0.02974                                       | 98.77%                        | Beta(1,1)        |
| 1000                               | 0.02382   | 0.02428                                       | 98.09%                        | Beta(1,1)        |

TABLE 1. Simulation results. Values are the arithmetic means over 30 trials. The functional is iterated 4 times starting with the uniform distribution on  $[0,1]$  as initial point. Functions are evaluated at 20 equally spaced points on  $[0,1]$ . The proposed estimator can be said to be almost equivalent as  $\hat{F}_n$ , the best estimator of  $F$ .

function  $h(t) = \lim_{\Delta t \rightarrow 0} P(t \leq T < t + \Delta t | T \geq t) / \Delta t = f(t) / S(t)$  and in particular the cumulative hazard function  $H(t) = \int_0^t h(s) ds = -\log S(t)$ . Usually  $T$  is thought to take values in  $[0, \infty)$ , but we can think to consider the estimation conditionally to the last sample failure, say  $\tau$ , and rescale the interval  $[0, \tau]$  to  $[0, 1]$ . So we will assume, from now on, all the failure times occur in  $[0, 1]$ , being 1 the instant of the last failure when the experiment stops. In this scheme of observation  $\hat{S}(t) = 1 - \hat{F}(t)$  is a natural estimator of  $S$ , with  $\hat{F}$  any estimator of  $F$  and, in particular, the IFS estimator. A more realistic situation is when some censoring occurs, in the sense that, as time pass by, some of the initial  $n$  observations are removed at random times  $C$  not because of failure (or death) but for some other reasons. In this case, a simple distribution function estimator is obviously not good. Let us denote by  $t_0 = 0 < t_1 < \dots < t_{d-1} < t_d = 1$  the observed instants of failure (or death). A well known estimator of  $S$  is the Kaplan-Meier estimator

$$\hat{S}(t) = \prod_{t_i < t} \frac{r(t_i) - d_i}{r(t_i)}$$

where  $r(t_i)$  are the subject exposed to risk of death at time  $t_i$  and  $d_i$  are the dead in the time interval  $[t_i, t_{i+1})$  (see the original paper of Kaplan and Meyer [8] or for a modern account [5]). In our case  $d_i$  is one as  $t_i$  are the instants when failures occur. Subjects exposed to risk are those still present in the experiment and not yet dead or censored. This estimator has good properties whenever  $T$  and  $C$  are independent. Related to the quantities  $r(t_i)$  and  $d_i$  it is also available the Nelson estimator for the function  $H$  that is defined as  $\hat{H}(t) = \sum_{t_i < t} d_i / r(t_i)$ . We assume for simplicity that there are not ties, in the sense that in each instant  $t_i$  only one failure occurs. The function  $\hat{H}(t)$  is a increasing step-function. Now let  $\hat{\mathcal{H}}(t) = \hat{H}(t) / \hat{H}(1)$ .  $\hat{\mathcal{H}}(t)$  can be thought as an empirical estimates of a distribution function  $\mathcal{H}$  on  $[0,1]$ . To derive an IFS estimator for the cumulative hazard function  $H$  we construct the sample quantiles by simply taking the inverse of  $\hat{\mathcal{H}}$ . Suppose we want to deal with  $k + 1$  quantiles, being  $\hat{q}_1 = 0$  and  $\hat{q}_{k+1} = 1$ . One possible definition of the empirical quantile of order  $m/k$  is obtained by the formula

$$\hat{q}_{m+1} = t_i + \frac{t_{i+1} - t_i}{\hat{\mathcal{H}}(t_{i+1}) - \hat{\mathcal{H}}(t_i)} \cdot \left( \frac{m}{k} - \hat{\mathcal{H}}(t_i) \right), \quad \text{if } \hat{\mathcal{H}}(t_i) \leq \frac{m}{k} < \hat{\mathcal{H}}(t_{i+1}), \quad (2)$$

for  $i = 0, 1, \dots, d - 1$  and  $m = 1, 2, \dots, k - 1$ . Now set  $p_i = 1/k$ ,  $i = 1, 2, \dots, k$  and  $\hat{q}_i$ ,  $i = 1, 2, \dots, k + 1$  as in (2). An IFS estimator of  $H$  is  $\hat{H}(1) \cdot \tilde{H}(t)$  where  $\tilde{H}(t)$  is the following IFS:

$$\tilde{H}(t) = \tilde{H}u(t) = \frac{1}{k}u\left(\frac{t - \hat{q}_i}{\hat{q}_{i+1} - \hat{q}_i}\right) + \frac{i}{k}, \quad t \in [\hat{q}_i, \hat{q}_{i+1}),$$

and  $u$  is any member of the space of distribution function on  $[0, 1]$ . In (2) we have assumed that  $\mathcal{H}$  is the distribution function of a continuous random variable, with  $\mathcal{H}$  varying linearly between  $t_i$  and  $t_{i+1}$ , but of course any other assumption than linearity can be made as well (for example an exponential behaviour). A Fleming-Harrington (or Altshuler) IFS-estimator of  $S$  is then

$$\tilde{S}(t) = \exp\{-\hat{H}(1) \cdot \tilde{H}(t)\}, \quad t \in [0, 1].$$

## 5. FINAL REMARKS ABOUT THE METHOD

There are several open issues in this topic as this is a first attempt to introduce IFS in distribution function estimation. The most important one is that a good characterization of the fixed point of the IFS is still missing. The second one, mostly not touched in the IFS literature, is how to choose the maps. We have suggested a quantile approach but some other good partition of the space, like a dyadic sequence, can be used at the cost of the need to solve some optimization problem. In [6] this problem is touched incidentally but not in a statistical perspective.

## REFERENCES

- [1] L. Arnold and H. Crauel, Iterated function systems and multiplicative ergodic theory, *Diffusion processes and related problems in analysis* (Pinsky M.A., Vihstutz V. eds.), vol. II, Stochastics flows, Progress in probability 27, Birkäuser, Boston, 283-305, 1992.
- [2] M.F. Barnsley and S. Demko, Iterated function systems and the global construction of fractals, *Proc. Roy. Soc. London, Ser A*, **399**(1985), 243-275.
- [3] C.A. Cabrelli, B. Forte, U.M. Molter and E.R. Vrscay, Iterated fuzzy set systems: a new approach to the inverse problem for fractals and other sets, *J. Math. Anal. Appl.*, **171**(1992), 79-100.
- [4] J.H. Elton and M. Piccioni, Iterated function systems arising from recursive estimation problems, *Probab. Theory Related Fields*, **91**(1992), 103-114.
- [5] T.R. Fleming and D.P. Harrington, Counting processes and survival analysis, Wiley, New York, 1991.
- [6] B. Forte and E.R. Vrscay, Solving the inverse problem for function/image approximation using iterated function systems, I. Theoretical basis, *Fractal*, **2**(1994), 3, 325-334.
- [7] J. Hutchinson, Fractals and self-similarity, *Indiana Univ. J. Math.*, **30**(1981), 5, 713-747.
- [8] E. Kaplan and P. Meyer, Nonparametric estimator from incomplete observations, *J. Amer. Statist. Ass.*, **53**(1958), 457-81.

- [9] A.A. Kwiecinska and W. Slomczynski, Random dynamical systems arising from iterated function systems with place-dependent probabilities, *Stat. and Prob. Letters*, **50**(2000), 401-407.
- [10] D. La Torre and M. Rocca, Iterated function systems and optimization problems, *Rend. Sem. Mat. Messina SER II*, **6**(1999), 21, 165-173.
- [11] P.W. Millar, Asymptotic minimax theorems for the sample distribution function, *Z. Wahrscheinlichkeitstheorie Verw. Geb.*, **48**(1979), 233-252.
- [12] P.W. Millar, The minimax principle in asymptotic statistical theory, *Lecture Notes in Math.*, **976**(1983), 76-265.
- [13] L. Montrucchio and F. Privileggi, Fractal steady states in stochastic optimal control models, *Ann. Op. Res.*, **88**(1999), 183-197.
- [14] R.T. Rockafellar and R.J-B. Wets, *Variational analysis*, Springer, New York, 1998.