The pvars R Package: VAR Modeling for Heterogeneous Panels

Lennart Empting
University of Duisburg-Essen

Abstract

pvars offers a seamless implementation of vector autoregressive (VAR) methods for heterogeneous panel data. The R package comprises *panel cointegration rank tests* which can account for *cross-sectional dependence* and for *structural breaks* in the deterministic terms. The implemented *panel SVAR models* can be estimated under these specifications with *pooled cointegrating vectors* and identified by various *panel identification procedures*. In this article, we review these methods and present their modular implementation in R. Two empirical illustrations reproduce examples from the literature step by step and guide the **pvars** users in conducting their own analyses.

Keywords: panel cointegration rank tests, panel vector autoregressive models, pooled cointegrating vectors, structural identification, independent components, R, **pvars**.

1. Introduction

The ever-increasing availability of macroeconomic data and recent developments in the multivariate time series analysis for panels have popularized panel vector autoregressive methods in applied econometric research. The **pvars** package summarizes a toolkit for the empirical analysis of panels in the narrow sense, where the same time series variables are repeatedly observed across several individual entities such as countries. This three-dimensional data structure thus contrasts with panels in a wider sense, where time series are just arranged along the same periods. Although **pvars**' application is not limited to financial or macroeconometric research, it addresses the data properties typically found in macroeconomic panels, namely (i) *mostly heterogeneous* (ii) *endogenous interactions* between (iii) potentially *nonstationary* variables. Since (iv) the *time dimension is distinctively larger* than the cross-section, the time series are prone to (v) a *complex deterministic term* with structural breaks in their mean and linear trend. The individual entities are usually subject to (vi) *cross-sectional dependence*.[1]

In order to deal with these data properties, the econometric literature extends individual time series methods by the cross-sectional dimension under selective pooling assumptions. For example, the panel methods implemented by the Stata commands `xtcointtest` (StataCorp 2025), `xtwest` (Persyn and Westerlund 2008), and `xtpmg` (Blackburne and Frank 2007)

[1]See Pedroni (2019) for an intuition and discussion of these panel properties and an overview of recent developments of single-equation and system-based methods. The data characteristics can be found in the applied econometrics of climate (Pretis 2020), energy (Smyth and Narayan 2015), and growth linked to financial development (Christopoulos and Tsionas 2004) or public capital (Empting and Herwartz 2025).

originate from the single-equation framework comprising residual-based cointegration tests and autoregressive distributed lag (ARDL) models in error-correction representation. In contrast, **pvars** relies on vector autoregressive (VAR) models as a system of equations. This approach has the advantage of avoiding restrictive exogeneity assumptions on the variables and modeling their dynamics and interactions explicitly. The model in vector error correction representation (VECM) can further accommodate multiple long-run relations and different types of deterministic regressors within the cointegration. Particularly the cointegration relations are reasonable candidates for a panel-wide pooling, while short-run dynamics are usually assumed to differ across individuals.

The synthesis of VAR models and methods for heterogeneous panels has not been implemented yet, but many packages for R (R Core Team 2026) can already contribute specification results and individual counter-checks. They can be subsumed into two pillars for **pvars**: First, the **vars** ecosystem with **urca** (Pfaff 2008a), **vars** (Pfaff 2008b), and **svars** (Lange, Dalheimer, Herwartz, and Maxand 2021) covers individual VAR methods such as VECM and structural identification. The second pillar is the single-equation panel methods from **plm** (Croissant and Millo 2008) such as dynamic panel regression models and panel unit root tests. Univariate panel time series could be tested for common or idiosyncratic nonstationarity using the more specialized package **PANICr** (Bronder 2016). Like **pvars**, the R package **panelvar** (Sigmund and Ferstl 2019) and its Stata equivalent **pvar** (Abrigo and Love 2016) rest on the two pillars. However, they focus on stationary panels that contain more individuals than periods ($N > T$), and the implemented estimators rely on slope coefficients that are homogeneous across all individuals in the PVAR model. Hence, they rather suit microeconomic applications, and their primary task is to deal with the *Nickell bias* (1981).[2]

The objective of **pvars** is to close the gap in the **vars** ecosystem and provide a seamless implementation of the fruitful panel VAR methodology. For this, three fields of VAR applications are integrated into **pvars**, namely panel estimation, cointegration testing, and structural identification. The implemented methods particularly noteworthy for each application field are (1) panel VECM with *pooled cointegrating vectors* from Breitung's (2005) two-step estimator. (2) The new panel cointegration rank tests by Arsova and Örsal (2017; 2018) respect cross-sectional dependence stemming from common factors. Arsova and Örsal (2021) consider also structural breaks in the deterministic term. (3) Data-driven identification methods based on *independent component analysis* (ICA) are extended for panels by Calhoun, Adali, Pearlson, and Pekar (2001) and Herwartz and Wang (2024).

The remainder of this article is structured as follows: Section 2 is a review of the econometric methods which are relevant for using this R package. Section 3 highlights the available functions in **pvars**, their implementation, and their library of auxiliary functions. In Section 4, we demonstrate two empirical applications of **pvars** to exemplary data from the reviewed articles. Section 5 summarizes this article. **pvars** can be directly installed from the *Comprehensive R Archive Network* (CRAN) and is shipped with documentation for further help.

2. Review of the econometric methodology

[2]See also Canova and Ciccarelli (2013) for a survey on panel VARs. Bayesian techniques are used in **BGVAR** (Boeck, Feldkircher, and Huber 2022) for Bayesian global VARs and in **bpvars** (Sanchez-Martinez and Woźniak 2026) for forecasting with Bayesian panel VARs.

A main motivation for combining multivariate time series from individual entities by panel methods is to increase the sample size and thereby extend the available information set for a more precise estimation and higher test power. Table 1 provides an overview of the literature of panel cointegration rank tests. In this unifying framework, the rows of the table classify the *pooling approach* according to the different stages of hypothesis testing, at which the cross-section information is presumed to be homogeneous. The listed pooling approaches allow for increasing heterogeneity between the individuals: (1) Panel-homogeneous cointegrating vectors enable a pooled estimation of these long-run coefficients. (2) The averaged statistic is standardized by moments of the distribution which the individual statistics have in common.[3] Finally, (3) the meta-analytical combination of individual p-values is the most flexible approach, where the commonality aims at a consistent test decision under an increasing number of individuals. If the alternative hypothesis is true, a non-vanishing share of individuals must actually exhibit this property.

Table 1: Panel tests for the rank of cointegration.

		First generation	**Second generation**		**Third generation**
	Cross-sectional dependence:	Independence	Correlated errors	PANIC (2004)	Correlated probits
Pooling approach:					
(1)	Pooled coefficients	Breitung (2005)	[a)] $LR(\Pi_C \mid \Pi_A)$	—	×
(2)	Averaged test statistics	Larsson et al. (2001) Örsal, Droge (2014)	[a)] $LR(\Pi_B \mid \Pi_A)$	Arsova, Örsal (2018) [b)] PMSB^{Z}	×
(3)	Meta-analytically combined p-values	Maddala, Wu (1999) Choi (2001)	—	Örsal, Arsova (2017) [b)] PMSB^{F}, PMSB^{C}	Hartung (1999) Arsova, Örsal (2021)

Not implemented in pvars: The panel tests by a) Groen, Kleibergen (2003) and b) Carrion-i-Silvestre, Surdeanu (2011).

The columns of Table 1 represent the *generations*, through which panel tests have evolved, and reflect the increasing complexity of the assumed data generating process. The first articles propose the plain approach for combining independent individual tests and thus are naive towards any dependencies between the individual entities. This cross-sectional dependence can emerge e.g. between countries due to their trade and financial relations and is a research topic itself with regard to the global business cycle, the crude oil price, or other important commodities of the world market. The prevalent data property decreases the actual information content in macroeconomic panels compared to cross-independent samples of same size. First-generation tests then reject a correct null hypothesis more often than the nominal significance level and are thus "over-sized". In contrast, panel tests of the second generation maintain a correct test size and third-generation tests are robust against structural breaks additionally. For cointegration rank tests, the econometric literature considers structural breaks in the mean or linear trend of the cointegration relations. Empirical examples which are associated with shifts and with breaks in the linear time trend are the German reunification, the beginning of the Great Moderation, and its discussed end in the wake of the Great Recession. For a proper data representation, the VAR model must respect those changes in the long-run equilibrium, towards which the error correction mechanism adjusts. Otherwise, the cointegration tests would fail to detect a reversion to the new equilibrium and

[3] In fact, the univariate counterpart by Im, Pesaran, and Shin (2003, p. 59, Remark 3.1) allows for individual-specific test distributions if their third moments exist. The approximating gamma distribution with its well-defined moments suggests that this holds true for cointegration rank tests. By the Lyapunov central limit theorem, the cross-sectional average of their first and second moments can then be used instead. In light of this, future studies may extend JMN (2000) and KN (2019) to panel tests with individual deterministic terms.

thus understate the cointegration rank.

A third dimension of panel test construction is constituted by the underlying individual procedures, which have not been mapped onto Table 1. These *branches* are heterogeneous and often used side-by-side in a single article such that they do not follow such a clear pattern as the other two panel test properties do. This does not only hold for the example of cointegration tests, but also for VAR estimation and structural identification. All panel applications are confronted with similar construction problems and thus follow these dimensions in principle. Since the individual methods elude any classification scheme for Table 1, Section 2.1 first explains their technical details and lays the foundation for Section 2.2 presenting the panel methods along the two dimensions of Table 1. The sections have the same structure according to estimation, testing, and identification as established in the individual and extended to the panel context respectively.

Notation. Subscript $t = 1, \dots, T$ denotes the periods and $k = 1, \dots, K$ refers to the endogenous variables in a multivariate time series. To simplify notation, Section 2.1 on the individual methods omits subscript $i = 1, \dots, N$ as the label specific to each of the N individuals. Section 2.2 uses i to distinguish individual-specific elements also from homogeneous elements. Without an individual identifier i, the latter remain the same over the complete cross-section. Moreover, symbol $\xrightarrow{d}$ designates convergence in distribution. The $(K \times (K-r))$ matrix $A_\perp$ is the orthogonal complement of a $(K \times r)$ matrix A such that $\text{rk}\{[A : A_\perp]\} = K$ indicates full rank and $(A_\perp)^\top A = 0$. By convention, the orthogonal complement of a nonsingular square matrix is the zero matrix 0 and the orthogonal complement of zero matrix 0 is an identity matrix I_K. With slight abuse of notation, integer $r_\perp$ refers to the number of stochastic trends in a multivariate time series, corresponding to the cointegration rank r.

2.1. Individual methods

The basic unit for all presented panel methods is an individual VAR model of the form

$$\boldsymbol{y}_t = \Phi \boldsymbol{d}_t + A_1 \boldsymbol{y}_{t-1} + ... + A_p \boldsymbol{y}_{t-p} + \boldsymbol{u}_t \quad \text{with} \quad \boldsymbol{u}_t \sim (0, \Sigma_u), \tag{1}$$

where $\boldsymbol{y}_t$ is a vector of K stacked time series, $A_j, j = 1, \dots, p$, are $K \times K$ coefficient matrices for the VAR process of order p, and Φ is a coefficient matrix for the deterministic regressors $\boldsymbol{d}_t$. The errors $\boldsymbol{u}_t$ are assumed to be serially, but not necessarily contemporaneously independent. Hence, the covariance matrix Σ_u is usually non-diagonal as the VAR model is estimated in reduced form initially. Finding the structural shocks $\boldsymbol{\epsilon}_t = \mathsf{B}^{-1}\boldsymbol{u}_t$, which exhibit no contemporaneous correlation, is the matter of structural identification in the decomposition problem $\Sigma_u = \mathsf{B}\mathsf{B}^\top$. The K variances of the structural shocks can be normalized to unity such that $\boldsymbol{\epsilon}_t \sim (0, I_K)$. Consequently, the unique identification of the impact matrix B requires at least $K(K-1)/2$ restrictions on the $K^2 - K$ covariances in Σ_u.

If $\boldsymbol{y}_t$ is integrated of order $I(1)$ at most and its first differences $\Delta\boldsymbol{y}_t$ are thus stationary, the VAR model of (1) in levels can be rewritten in its vector error correction representation

$$\begin{aligned} \Delta\boldsymbol{y}_t \;&=\; \Phi\boldsymbol{d}_t + \Pi_K \boldsymbol{y}_{t-1} + \sum_{j=1}^{p-1} \Gamma_j \Delta\boldsymbol{y}_{t-j} + \boldsymbol{u}_t \quad \text{with} \quad \boldsymbol{u}_t \sim (0, \Sigma_u) \\ \text{using} \quad \Pi_K &:= -I_K + \sum_{j=1}^{p} A_j \quad \text{and} \quad \Gamma_j := -\sum_{j^*=j+1}^{p} A_{j^*},\ j = 1, \dots, p-1. \end{aligned} \tag{2}$$

The $K \times K$ coefficient matrix $\Pi_K = \alpha\beta_K^\top$ of the error correction term summarizes the $K \times r$ cointegrating matrix β_K and loading matrix α, which adjusts the disequilibria in the r long-run cointegration relations $\beta_K^\top \boldsymbol{y}_{t-1}$. The regressors $\boldsymbol{d}_t$ of the deterministic term $\Phi\boldsymbol{d}_t = [\Phi_1 : \Phi_2]\left(\boldsymbol{d}_{1,t}^\top, \boldsymbol{d}_{2,t}^\top\right)^\top$ are either unrestricted ($\boldsymbol{d}_{2,t}$) or restricted ($\boldsymbol{d}_{1,t}$) to the cointegration relations such that the coefficients Φ_1 emerge from $\Pi := [\Pi_K : \Phi_1] = \alpha\left[\beta_K^\top : \beta_0^\top\right] = \alpha\beta^\top$. Finally, each Γ_j is a $K \times K$ matrix of coefficients for short-run effects at lag $j = 1, \ldots, p-1$.

Estimating cointegrated VAR models

Johansen (1991, 1996) provides a comprehensive collection of *maximum likelihood* (ML) methods for the individual VECM in (2). His ML estimators and LR-tests are the foundation for any cointegrated VAR method of the **pvars** package. Hence, their definitions and components are throughout used in this article and shall be introduced here briefly. For convenience, first rewrite (2) in compact notation as

$$Z_0 \;= \alpha\beta^\top Z_1 + \Gamma Z_2 + U \quad \text{now with} \quad \boldsymbol{u}_t \sim \mathcal{N}(0, \Sigma_u), \tag{3}$$

where the matrix $\Gamma := [\Gamma_1 : \cdots : \Gamma_{p-1} : \Phi_2]$ lines up the coefficients for the short-run dynamics and for the unrestricted deterministic regressors. The observed time series are collected in the $K \times T$ matrix $Y := [\boldsymbol{y}_1, \ldots, \boldsymbol{y}_T]$ for a sample $1, \ldots, T$ and the same holds for the error matrix $U := [\boldsymbol{u}_1, \ldots, \boldsymbol{u}_T]$, which contains no presample periods by construction of the estimation with lagging regressors. Accordingly, the regressand matrix is given by $Z_0 := \Delta Y$ and the two regressor matrices with T columns alike by

$$Z_1 := \begin{bmatrix} \boldsymbol{y}_0 & \cdots & \boldsymbol{y}_{T-1} \\ \boldsymbol{d}_{1,1} & \cdots & \boldsymbol{d}_{1,T} \end{bmatrix} \quad \text{and} \quad Z_2 := \begin{bmatrix} \Delta\boldsymbol{y}_0 & \cdots & \Delta\boldsymbol{y}_{T-1} \\ & \vdots & \\ \Delta\boldsymbol{y}_{1-p+1} & \cdots & \Delta\boldsymbol{y}_{T-p+1} \\ \boldsymbol{d}_{2,1} & \cdots & \boldsymbol{d}_{2,T} \end{bmatrix}. \tag{4}$$

The variables for the error correction term enter Z_1 in levels. Z_2 stacks the first-differenced and the unrestricted deterministic regressors. If $p = 1$, all lagged $\Delta\boldsymbol{y}_t$ drop out of Z_2. In conditional VECM, vectors of L (weakly) exogenous variables $\boldsymbol{x}_t$ could be included as lagged $I(1)$ regressors into Z_1 as well as instantaneous and up to q lagged short-run effects into Z_2. The estimation would follow the same proceeding (Pesaran, Shin, and Smith 2000; Lütkepohl 2005, p. 398), but we skip these terms here for the sake of notational brevity.

ML estimation.[4] The first order conditions from the maximized log-likelihood function

$$\begin{aligned} \ln\mathcal{L}(\alpha, \beta, \Gamma, \Sigma_u) = &-\frac{KT}{2}\ln(2\pi) - \frac{T}{2}\ln\left(\det(\Sigma_u)\right) \\ &- \frac{1}{2}\mathrm{tr}\left[\left(Z_0 - \alpha\beta^\top Z_1 - \Gamma Z_2\right)^\top \Sigma_u^{-1} \left(Z_0 - \alpha\beta^\top Z_1 - \Gamma Z_2\right)\right] \end{aligned} \tag{5}$$

are solved for the estimators of the VECM, which corresponds to the following three steps:

1. *Concentrate out* short-run effects and the unrestricted deterministic term. The first step of Π's estimation is to remove the component $\Gamma\boldsymbol{z}_{2t}$ from $\Delta\boldsymbol{y}_t$ and $\boldsymbol{y}_t$ by OLS-regression of Z_0 resp. Z_1 on Z_2. Using the Frisch-Waugh Theorem, the projection

[4] See Johansen (1996, Ch. 6), Lütkepohl (2005, Ch. 7.2.3) and Lütkepohl (2006, Ch. 3.1) for a more detailed derivation and explanation.

matrix $M_2 := I_T - Z_2^\top (Z_2 Z_2^\top)^{-1} Z_2$ generates the "long-run residuals" $R_0 = Z_0 M_2$ resp. $R_1 = Z_1 M_2$ such that (3) collapses into the plain error correction model

$$\begin{aligned} Z_0 M_2 &= \alpha\beta^\top Z_1 M_2 + \Gamma Z_2 M_2 + U M_2, \\ R_0 &= \alpha\beta^\top R_1 + \widetilde{U}. \end{aligned} \tag{6}$$

2. Estimate the concentrated model in (6) by *reduced-rank regression* (RRR) according to Anderson (1951).[5] For doing so, define the moment matrices

$$S_{00} := \frac{R_0 R_0^\top}{T}, \quad S_{01} := \frac{R_0 R_1^\top}{T}, \quad \text{and} \quad S_{11} := \frac{R_1 R_1^\top}{T}. \tag{7}$$

Further inserting the OLS estimator from (9) for $\widetilde{\alpha}(\beta)$ into the concentrated likelihood function transforms the ML maximization into the generalized eigenvalue problem

$$\det\left(\lambda S_{11} - S_{01}^\top S_{00}^{-1} S_{01}\right) = 0. \tag{8}$$

This equation has K solutions with the ordered eigenvalues $\lambda_1 \geq \lambda_2 \geq \ldots \geq \lambda_K \geq 0$, which are the squared *canonical correlation coefficients* and indicate the "strength" of cointegration in each solution. By the definition of cointegration, the correct coefficients $\alpha\beta^\top$ must recombine the nonstationary R_1 to the stationary R_0. Hence, the estimator $\widetilde{\beta}$ equals the eigenvectors of the r largest λ_j and is superconsistent converging with rate T instead of the usual $\sqrt{T}$ of the remaining OLS estimates. The $r_\perp$ eigenvectors with the smallest λ_j are associated with the underlying stochastic trends and thus must cancel out from the henceforth *rank-restricted* VECM.

3. Maximize the log-likelihood function in (5) conditional on a given β, for which the superconsistent RRR-estimate $\widetilde{\beta}$ can be inserted. The solutions for the remaining estimators of the reduced-form VECM in (3) are in fact *conditional OLS*, namely

$$\begin{aligned} \widetilde{\Pi} &= \widetilde{\alpha}\beta^\top = \left[R_0 \left(\beta^\top R_1\right)^\top \left(\left(\beta^\top R_1\right)\left(\beta^\top R_1\right)^\top\right)^{-1}\right]\beta^\top = S_{01}\beta\left(\beta^\top S_{11}\beta\right)^{-1}\beta^\top, \\ \widetilde{\Gamma} &= \left(Z_0 - \widetilde{\Pi} Z_1\right) Z_2^\top (Z_2 Z_2^\top)^{-1}, \\ \widetilde{U} &= Z_0 - \widetilde{\Pi} Z_1 - \widetilde{\Gamma} Z_2 = R_0 - \widetilde{\Pi} R_1. \end{aligned} \tag{9}$$

The residual covariance matrix is given by $\widetilde{\Sigma}_u = \frac{\widetilde{U}\widetilde{U}^\top}{T} = S_{00} - \widetilde{\Pi} S_{01}^\top$ for ML estimation and is further corrected for n, the number of coefficients per equation, in the OLS estimator $\widehat{\Sigma}_u = \frac{T}{T-n} \cdot \widetilde{\Sigma}_u$. From (9), it becomes clear that only the loadings $\widetilde{\alpha}(\beta)$ adjust towards the normalization of β, while all other estimators are conditional on the complete matrix product $\widetilde{\Pi} = \widetilde{\alpha}\beta^\top$. This shows that the cointegrated VAR process in (2) and its impulse response functions (IRF) are invariant to any eligible normalization under a chosen rank r of Π.

GLS-based trend adjustment. For the unique purpose of determining the cointegration rank, Lütkepohl and Saikkonen (2000) suggest estimating and removing the deterministic term prior to an LM-test. Saikkonen and Lütkepohl (2000c) apply the LR-statistic in (15) to the "detrended" time series, which may further improve the test power. This *SL-procedure*

[5] See also Anderson (2003, Ch. 12.7) and Izenman (1975).

entails a series of publications proposing different specifications of the deterministic term, which can be subsumed by the additive data generating process

$$\begin{aligned} \boldsymbol{y}_t &= M_\mu \boldsymbol{d}_t + \boldsymbol{y}_t^{\not d}, \\ \boldsymbol{y}_t^{\not d} &= A_1 \boldsymbol{y}_{t-1}^{\not d} + \ldots + A_p \boldsymbol{y}_{t-p}^{\not d} + \boldsymbol{u}_t^{\not d}. \end{aligned} \tag{10}$$

Therein, the VAR process of the "pure" stochastic component $\boldsymbol{y}_t^{\not d}$ is usually abbreviated by a lag polynomial $A(L) = I_K - A_1 L - \ldots - A_p L^p$. The $K \times n_d$ matrix M_μ collects the coefficients of the deterministic term in the moving-average representation of the VAR. Specifications of $\boldsymbol{d}_t$ pursuant to the SL literature are listed in Table 2 and illustrated in Section 4.2 in detail.

All SL-procedures consider a *GLS estimator* to determine M_μ. It becomes feasible by a preceding ML estimation of the VECM in (2), for which Table 2 indicates also the conforming deterministic cases. Each hypothesis $r_{H0} = 0, \ldots, K-1$ in the test sequence then requires its specific rank restriction on the ML estimates and an own GLS estimation of M_μ. For this, the ML estimates of the rank-restricted VECM are converted into $\widetilde{A}_1, \ldots, \widetilde{A}_p$ of the VAR in levels and inserted into the transformed model

$$Q^\top A(L) \boldsymbol{y}_t = Q^\top A(L) M_\mu \boldsymbol{d}_t + Q^\top \boldsymbol{u}_t^{\not d} \quad \text{for} \quad t = 1, \ldots, T. \tag{11}$$

Therein, $\boldsymbol{d}_t$ and $\boldsymbol{y}_t$ are set to initial zeros in any presample period $t \leq 0$ and the feasible transformation matrix $\widetilde{Q}$ is obtainable by inserting ML estimates into

$$Q = \left[\Sigma_u^{-1} \alpha \left(\alpha^\top \Sigma_u^{-1} \alpha \right)^{-1/2} : \alpha_\perp \left((\alpha_\perp)^\top \Sigma_u \alpha_\perp \right)^{-1/2} \right]. \tag{12}$$

The left-multiplication of $Q^\top A(L)$ to (10) in (11) allows for subtracting the confounding effects of the VAR dynamics and transforms the residual covariance matrix into I_K. After vectorizing the model in (11), M_μ can thus be estimated by OLS. Alternatively, Saikkonen and Lütkepohl (2000c, p. 438) show that

$$QQ^\top = \Sigma_u^{-1} \alpha \left(\alpha^\top \Sigma_u^{-1} \alpha \right)^{-1} \alpha^\top \Sigma_u^{-1} + \alpha_\perp \left((\alpha_\perp)^\top \Sigma_u \alpha_\perp \right)^{-1} (\alpha_\perp)^\top = \Sigma_u^{-1}. \tag{13}$$

Accordingly, the GLS estimator based on the estimated covariance matrix $\widetilde{\Sigma}_u$ can be applied directly to the data unaltered by $\widetilde{Q}$, but still corrected for $\widetilde{A}(L)$. Both proceedings lead to identical estimation results $\widehat{M}_\mu$. Finally, the deterministic term is subtracted and the VECM of the stochastic component $\hat{\boldsymbol{y}}_t^{\not d} = \boldsymbol{y}_t - \widehat{M}_\mu \boldsymbol{d}_t$ is estimated by ML again – now without any deterministic term. The usual LR-statistic in (15) can assess the sole null hypothesis of r_{H0}, but its test distribution in (18) differs from the Johansen procedure.

Deterministic term. Johansen and Nielsen (2018) distinguish between the *innovative* and the *additive* formulation of the deterministic term in (2) for the Johansen procedure resp. in (10) for the SL-procedure. Table 2 adopts these labels and lists the model specifications. Like in a stable VAR process, the deterministic regressors $\boldsymbol{d}_t$ in the VECM of (2) can contain either no deterministic component, an intercept, or an additional linear trend. Beyond these "standard" types, nonstationarity and cointegration do complicate the estimation and testing in the presence of deterministic terms. For example, if a constant is assigned to $\boldsymbol{d}_{2,t}$, the unit roots in the VECM do not only accumulate the innovations $\boldsymbol{u}_t$ into stochastic trends, but also this deterministic constant into a linear trend. On the other hand, the deterministic

Table 2: Verified specifications of the deterministic term.

Deterministic regressors		**Type**		**Literature**
restricted $\boldsymbol{d}_{1,t}$	unrestricted $\boldsymbol{d}_{2,t}$	innovative[a)]	additive[b)]	
• ***conventional:***				
none	none	*Case 1*	(needless)	—
constant	none	*Case 2*	*SL_mean*	L&S (2000, p. 185)
none	constant	*Case 3*	*SL_ortho*	S&L (2000a) is irrelevant for **pvars**
linear trend	constant	*Case 4*	*SL_trend*	L&S (2000)
none	constant & linear trend	*Case 5*	—	—
• ***period-specific:***				
none	seasonal dummies	+ *Case 2–5*	+ all	Johansen (1996, p. 84) || ...
impulse dummy	none	+ all	+ all	... TSL (2008, p. 348)
shift dummy	impulse dummies	+ *Case 2* [c)]	+ *SL_mean, SL_trend*	JMN (2000, Ch. 3.2) || S&L (2000b)[d)]
trend break	shift & impulse dummies	+ *Case 4*	+ *SL_trend*	JMN (2000, Ch. 3.1) || TSL (2008)

[a)] Case labeling in accordance with Juselius (2006, Ch. 6.3) and Hlouskova and Wagner (2010, p. 193).
[b)] Overview of SL-test specifications from Trenkler (2008, p. 24, Tab. 1; p. 25, Tab. 2).
[c)] For shifts in *Case 4*, Johansen *et al.* (JMN, 2000, Ch. 4) assess r_{H0} under trend breaks and then restrictions on β_0.
[d)] Lütkepohl, Saikkonen, and Trenkler (TSL, 2004) propose estimators of the unknown shift periods $\boldsymbol{\tau}$.

regressors $\boldsymbol{d}_{1,t}$ are restricted to the stationary cointegration relation and, thus, a linear trend in $\boldsymbol{d}_{1,t}$ does not generate a quadratic trend in the data. In the additive formulation of the SL-procedure, the stochastic component contains the unit roots so that they do not interfere with the deterministic component. Irrespective of the accumulating innovations, the deterministic term thereby retains its intended specification, as it is visible in the data, and can be easily subtracted after FGLS estimation. For this, each row in Table 2 juxtaposes the specifications of the additive model in (10) and the corresponding term in the innovative VECM of (2) which provides the parameters in (11)–(13) for feasible GLS estimation.

Additionally to the conventional types, there can be period-specific shifts in the mean and breaks in the linear trend as well as impulse dummies for a single period or a repeated pattern of dummies for deterministic seasonality. Both test procedures retain the conventional asymptotics in (19) if solitary impulse dummies are added or if the seasonal dummies in $\boldsymbol{d}_{2,t}$ are centered around zero and thus accumulate along a constant. In contrast, only the distributions $Z_{r_\perp}$ of the SL-procedure are invariant to the inclusion of shift dummies irrespective of their known or estimated shift period and both procedures must cope with the nuisance introduced by broken trend slopes. Johansen, Mosconi, and Nielsen (JMN, 2000), Kurita and Nielsen (KN, 2019) as well as Trenkler, Saikkonen, and Lütkepohl (TSL, 2008) provide solutions for the Johansen- and SL-procedures, respectively. Beyond these specifications whose asymptotics are verified by the literature of Table 2 and described in Section 2.1.2, **pvars** accepts all period-specific extensions technically and does not check their validity for the cointegration tests.

Like for the conventional types in Table 2, any regressor $\boldsymbol{d}_{1,t}$ in the cointegration term has a first-differenced counterpart in $\boldsymbol{d}_{2,t}$. While the additive model separates clearly between deterministic and stochastic components, the innovative VECM in (2) mixes the structural breaks into the autoregressive dynamics. Hence, the innovative model implies not only a single, but additional lagged first differences of the break over the periods $\tau, \tau+1, \ldots, \tau+(p-1)$ after the break occurring in period τ. Their coefficients depend on the other model parameters and would require nonlinear estimation. Against this, all authors of the reviewed literature accept a minor loss in degrees of freedom and prefer to estimate these coefficients separately without those restrictions. Like TSL (2008, p. 335), **pvars** sets lagged impulse dummies for

$\tau, \ldots, \tau + p - 1$ in $\boldsymbol{d}_{2,t}$ even after a trend break because, in combination with the shift in τ, they control for these nonlinear dynamics the same way as lagged shift dummies. In return, a subsequent FGLS estimation of M_μ in the additive (10) uses the regressor matrix $\boldsymbol{d}_t$ without these lagged first differences. Besides the conventional term, it contains only the trend break and shift at the very same period τ as illustrated in (38).

Testing the cointegration rank

In order to determine the cointegration rank r in the K-dimensional system of (2), all testing procedures imply a sequential decision making according to the hypotheses

$$H_0 : \operatorname{rk}(\Pi) = r_{H0} \quad \text{versus} \quad H_1 : \operatorname{rk}(\Pi) > r_{H0}, \quad r_{H0} = 0, \ldots, K-1. \tag{14}$$

As long as a null hypothesis is rejected, r_{H0} is increased by 1 and tested again. The procedure stops at the first non-rejected $r = r_{H0}$ or if the maximal $r_{H0} = K - 1$ has been rejected in favor of $r = K$. Correspondingly, the number of stochastic trends $r_\perp = K - r_{H0}$ declines in each step from K down to 1. Note that, under a given r_{H0}, the *maximum eigenvalue test* builds on the H_1 of $\operatorname{rk}(\Pi) = r_{H0} + 1$ and the *trace test* on $\operatorname{rk}(\Pi) = K$. Although both tests are applicable to panel extensions, only variants of the trace test have been adopted by the panel literature and are thus introduced here.

LR-test. Johansen (1988) developed the now predominant *likelihood ratio* test which compares the maximized likelihood $\mathcal{L}(r_{H0})$ from the rank-restricted model against the likelihood $\mathcal{L}(K)$ from the full-rank model with K variables. This trace statistic is defined as

$$\begin{aligned} \lambda^{\mathrm{LR}}(r_{H0}) &= -2\left[\ln \mathcal{L}(r_{H0}) - \ln \mathcal{L}(K)\right] \\ &= -T \sum_{j=r_{H0}+1}^{K} \ln(1 - \hat{\lambda}_j) \xrightarrow{d} Z_{r_\perp}. \end{aligned} \tag{15}$$

The estimated eigenvalues $\hat{\boldsymbol{\lambda}}$ are the squared canonical correlation coefficients obtained from the RRR in Section 2.1.1, and the number of nonzero eigenvalues λ_k equals the rank of Π. Correspondingly, this test assesses whether any of the $r_\perp$ smallest eigenvalues is significantly different from zero and thus increases the rank of Π. Under H_0, the asymptotic test distribution is a function of $r_\perp$-dimensional Brownian motions as outlined in (18).

LM-test. Despite its marginal role in the individual time series methodology, the *Lagrange multiplier* test adopted by Luukkonen, Ripatti, and Saikkonen (1999) has proven useful for the panel test procedure proposed by Breitung (2005) as it can serve as a vehicle to introduce the panel-homogeneous cointegrating vectors into an individual testing procedure. In order to make the cointegration rank r_{H0} testable by LM, the concentrated model in (6) is extended by $\phi(\beta_\perp)^\top R_1$ and multiplied by $(\alpha_\perp)^\top$. Consequently in the auxiliary model

$$\begin{aligned} R_0 &= \alpha\beta^\top R_1 + \phi(\beta_\perp)^\top R_1 + U^{LM} \\ \text{resp.} \quad (\alpha_\perp)^\top R_0 &= \phi^*(\beta_\perp)^\top R_1 + (\alpha_\perp)^\top U^{LM}, \end{aligned} \tag{16}$$

the extension is $H_0 : (\alpha_\perp)^\top \phi = \phi^* = 0_{r_\perp \times r_\perp}$ under r_{H0}, but converts $\Pi = \alpha\beta^\top + \phi(\beta_\perp)^\top$ into a full-rank matrix if the data suggest the alternative $H_1 : \operatorname{rk}(\Pi) = K$. After inserting suitable estimates for the orthogonal complements,[6] the coefficient matrix ϕ^* is estimated by OLS

[6]For his panel cointegration test, Breitung (2005) uses the individual first-step estimate for α_i and the pooled second-step estimate for β_K under rank restriction r_{H0}. In **pvars**, their orthogonal complements are then calculated via the QR-decomposition as done in the **MASS** package (Venables and Ripley 2002).

with regressand $E := \widehat{\alpha_\perp}^\top R_0$ and regressor $W := \widehat{\beta_\perp}^\top R_1$. The test statistic for $H_0 : \phi^* = 0$ is constructed according to the here all-equivalent LR-, Wald-, or LM-principle and is thus directly calculated with

$$\lambda^{\mathrm{LM}}(r_{H0}) = T \operatorname{tr}\left[EW^\top \left(WW^\top\right)^{-1} WE^\top \left(EE^\top\right)^{-1}\right] \xrightarrow{d} Z_{r_\perp}. \tag{17}$$

This LM-test statistic and Johansen's LR-test statistic from (15) are asymptotically identical and so are their test distributions.

Asymptotic distribution.[7] Under H_0 and $T \to \infty$, the test distribution converges to

$$Z_{r_\perp} = \operatorname{tr}\left[\int_0^1 \left(\mathrm{d}W^\circ_{r_\perp}\right) W^{\bullet\top}_{r_\perp} \left(\int_0^1 W^\bullet_{r_\perp} W^{\bullet\top}_{r_\perp}\right)^{-1} \int_0^1 W^\bullet_{r_\perp} \left(\mathrm{d}W^\circ_{r_\perp}\right)^\top\right] \tag{18}$$

$$\text{with} \quad W^\bullet_{r_\perp} := \begin{cases} W_{r_\perp}(s) & \text{for } \textit{Case 1} \text{ or } \textit{SL_mean} \\ W_{r_\perp}(s) - \int_0^1 W_{r_\perp}(s)\mathrm{d}s & \text{for } \textit{Case 2} \\ \left[W_{r_\perp}{}_s^{-1}(s)\right] - \int_0^1 \left[W_{r_\perp}{}_s^{-1}(s)\right] \mathrm{d}s & \text{for } \textit{Case 3} \\ W_{r_\perp}(s) - sW_{r_\perp}(1) & \text{for } \textit{Case 4} \text{ or } \textit{SL_trend} \end{cases} \tag{19}$$

$$\text{and} \quad \mathrm{d}W^\circ_{r_\perp} := \begin{cases} \mathrm{d}W_{r_\perp}(s) & \text{for } \textit{SL_mean} \text{ or any } \textit{Case} \\ \mathrm{d}W_{r_\perp}(s) - \mathrm{d}sW_{r_\perp}(1) & \text{for } \textit{SL_trend}. \end{cases}$$

The $r_\perp$-dimensional vector $W_{r_\perp}(s)$ stacks the independent standard Brownian motions. In (18), their vector products are integrated over their complete domain $s \in [0, 1]$. If the VECM accommodates deterministic regressors, $W^\bullet_{r_\perp}(s)$ must be specified accordingly. (19) lists the Brownian motions resp. bridges for the conventional cases from Table 2. In contrast to those, trend breaks are specific to the periods of their occurrence and the relative position of these periods within the sample affects $Z_{r_\perp}$ additionally. Also note that any unrestricted deterministic term which is accumulated under $I(1)$, e.g. the constant in *Case 3*, dominates the stochastic trends and thus replaces a Brownian motion in $W_{r_\perp}(s)$.

Simulated distribution. Since the asymptotic distribution of $Z_{r_\perp}$ is non-standard, critical values have been simulated e.g. by Osterwald-Lenum (1992). For this, $W_{r_\perp}$ is substituted by repeated simulations of $r_\perp$-dimensional vectors of Gaussian random walks with sufficiently[8] large sample size. Further, $Z_{r_\perp}$ can be approximated by the gamma distribution $\Gamma(\mathsf{s}, \mathsf{r})$, whose continuous probability density function offers the advantage of retrieving p-values conveniently. In order to do so, its shape- and rate-parameters are equated with

$$\mathsf{s} = \frac{\mathbb{E}(Z_{r_\perp})^2}{\mathrm{Var}(Z_{r_\perp})} \qquad \text{and} \qquad \mathsf{r} = \frac{\mathbb{E}(Z_{r_\perp})}{\mathrm{Var}(Z_{r_\perp})}. \tag{20}$$

Simulated values for the moments $\mathbb{E}(Z_{r_\perp})$ and $\mathrm{Var}(Z_{r_\perp})$ can be found for example in Larsson *et al.* (2001) and Breitung (2005).[9] In general, the simulation of $Z_{r_\perp}$ to retrieve the moments

[7] See Johansen (1996, Ch. 11.2), Lütkepohl (2005, Ch. 8.2), and Trenkler (2008, p. 24, Eq. 2.9).

[8] For example, Johansen (1996, Ch. 15) recommends $T = 400$. Breitung (2005, p. 171, App. B) employs $T = 500$ and Örsal and Droge (2014) $T = 1000$ in order to simulate the respective moments of $Z_{r_\perp}$.

[9] Larsson *et al.* (2001) provide moments of $Z_{r_\perp}$ only for *Case 1* and Breitung (2005) for *Case 2* to *Case 4* in order to center and scale the LR-bar panel test statistic as in (28). In **pvars**, these moments are stored in the internal list object `pvars:::coint_moments`.

or critical values depends on *(i)* the number of stochastic trends $r_\perp$ under H_0, *(ii)* the deterministic term, and *(iii)* the number of weakly exogenous variables in a conditional VECM. Hence, the different test specifications imply a grid of simulation setups to run.

More flexibility towards adapting these specifications is offered by response surface approximation of the moments as employed by Doornik (1998). For each *Case*, he estimates response surface coefficients[10] of a polynomial regression model $f(r_\perp)$ which "predicts" the moments of $Z_{r_\perp}$ for Johansen (1996). Trenkler (2008) derives response surface coefficients specifically for the trend-adjusted tests of (10). TSL (2008) tabulate these coefficients for trend-adjusted tests which accommodate structural trend breaks in the cointegration relationship. Likewise, JMN (2000) consider structural breaks in the constant and in the linear trend of the innovative model, which KN (2019) generalize for conditional VECM by Doornik (1998, Ch. 9). Although both models, additive and innovative, can cope with several breaks in principle, the regression models $f\left(r_\perp, \frac{\tau}{T}\right)$ by JMN (2000), KN (2019), and TSL (2008) contain polynomials of up to two break periods $\boldsymbol{\tau}$ only. **pvars** resorts to the respective approximation automatically in order to comply with the test specification selected by the user.

Identifying structure

Based on the given rank-restriction r and structural shocks $\boldsymbol{\epsilon}_t = \mathsf{B}^{-1}\boldsymbol{u}_t$, the VECM in (2) can be transformed according to the *Granger representation theorem* (GRT) into

$$\begin{aligned} \boldsymbol{y}_t &= \Xi\mathsf{B}\sum_{j=1}^{t}\boldsymbol{\epsilon}_j + \sum_{j=0}^{\infty}\Xi_j^*\mathsf{B}\boldsymbol{\epsilon}_{t-j} + \Xi\Phi\sum_{j=1}^{t}\boldsymbol{d}_j + \sum_{j=0}^{\infty}\Xi_j^*\Phi\boldsymbol{d}_{t-j} + \boldsymbol{y}_0^* \quad \text{with} \\ \Xi &= \beta_\perp\left[(\alpha_\perp)^\top\left(I_K - \sum_{j=1}^{p-1}\Gamma_j\right)\beta_\perp\right]^{-1}(\alpha_\perp)^\top. \end{aligned} \tag{21}$$

The initial vector $\boldsymbol{y}_0^*$ and the deterministic terms are usually dropped in order to trace the responses to an isolated structural impulse ϵ_{kt}. The number of stochastic trends $\mathrm{rk}(\Xi) = K - r = r_\perp$ follows from the $K \times r_\perp$ orthogonal complements $\alpha_\perp$ and $\beta_\perp$ for the $K \times r$ loadings α and cointegrating vectors β_K. Their different specifications illustrate the role of the components in (21) and their options for structural restrictions:

- If $r_\perp = 0$, the stochastic trends cancel out and the sole multiplier matrices $\Xi_j^*\mathsf{B}$ reflect the IRF of the implied *stable* SVAR model. Hence, just-identification requires the $K(K-1)/2$ restrictions to be imposed on the instantaneous effects B exclusively. A simple example is Sims (1980), who assumes recursive (short-run) causality such that the structural effects on impact can be calculated by $\widehat{\mathsf{B}} = \mathrm{chol}\left(\widehat{\Sigma}_u\right)$.
- If $r_\perp = K$, the VECM loses its error correction term and is thus estimated as a first-differenced VAR model by OLS. Since $\alpha_\perp = \beta_\perp = I_K$, matrix $\Xi = \left(I_K - \sum_{j=1}^{p-1}\Gamma_j\right)^{-1} = \Gamma(1)^{-1}$ is full-rank and $K(K-1)/2$ long-run restrictions are sufficient for identifying this SVAR of *growth rates* $\Delta\boldsymbol{y}_t$. In this example, recursive long-run causality is imposed on the structural long-run effects $\Xi\mathsf{B}$ such that $\widehat{\mathsf{B}} = \widetilde{\Gamma}(1)\ \mathrm{chol}\left(\widehat{\Sigma}_\infty\right)$ is calculated via the long-run covariance $\Sigma_\infty := \Xi\mathsf{B}\mathsf{B}^\top\Xi^\top = \Gamma(1)^{-1}\Sigma_u\left(\Gamma(1)^{-1}\right)^\top$.

[10] In **pvars**, the response surface coefficients are stored in the internal list object `pvars:::coint_rscoef`.

- A cointegration rank r implies $r_\perp$ permanent shocks at minimum and conversely r transitory shocks at maximum. Identification schemes in the tradition of King, Plosser, Stock, and Watson (1991) assume that $\Xi\mathsf{B}$ is rank-deficient due to r columns of $\mathbf{0}_K$. Each $\mathbf{0}_K$ suppresses the stochastic trends and thus defines a transitory shock in $\boldsymbol{\epsilon}_t$. Other formulations of the rank-deficiency may lead to fewer transitory shocks.

ML estimation.[11] The impact matrix B can be estimated by ML via $\mathcal{L}(\alpha, \beta, \Gamma, \mathsf{B})$ as an extension of (5). Inserting the reduced-form estimates from Section 2.1.1 facilitates a conditional ML estimation of B as Step **4.** with the concentrated likelihood function

$$\ln \mathcal{L}_c(\mathsf{B}) = -\frac{KT}{2}\ln(2\pi) - \frac{T}{2}\ln\left(\det(\mathsf{B})\right)^2 - \frac{1}{2}\mathrm{tr}\left[\mathsf{B}^{\top-1}\mathsf{B}^{-1}\widetilde{\Sigma}_u\right]. \tag{22}$$

This constrained maximization problem has no closed-form solution. Hence, Breitung, Brüggemann, and Lütkepohl (2004) maximize $\mathcal{L}_c(\mathsf{B})$ with respect to the free parameters subject to the identifying restrictions numerically by the scoring algorithm of Amisano and Giannini (1997). If the number of short- and long-run restrictions suffices only for just-identification, the equality $\widetilde{\Sigma}_u = \widetilde{\mathsf{B}}\widetilde{\mathsf{B}}^\top$ holds and the scoring algorithm serves as a nonlinear equation solver.

2.2. Panel methods

Estimating cointegrated VAR models

In macroeconomic panels, the idiosyncrasies of countries can subvert the objective to increase estimates' precision with pooled data. Deterministic effects like intercepts are usually kept individual-specific in panel data analysis, but even the assumption of homogeneous slope coefficients can be too restrictive. Pesaran and Smith (1995) demonstrate how systematic differences between individual dynamics lead to inconsistent estimates and consequently deny poolability. As an economically plausible middle ground, Pesaran, Shin, and Smith (1999) propose a selective pooling for single-equation models, where only the long-run relationship is homogeneous across all individuals, while the short-run dynamics are considered as heterogeneous. The following panel estimators adopt this principle into a system of equations and restrict the r cointegrating vectors $\beta_i = \beta \;\forall\; i$. Although the individual VECM in (2) does not require the basic normalization of the error correction term as single-equation models do, a panel-wide normalization becomes necessary in order to extract the homogeneous β from the individual coefficient matrices $\Pi_i = \alpha_i\beta^\top$.

Two-step estimator.[12] Breitung (2005) extends the individual two-step estimator by Ahn and Reinsel (1990) to a panel estimator for β. Based on the panel-wide normalization $\Pi_i = \alpha_i\left[I_r : \mathbf{B}\right] = \left[\alpha_i I_r : \alpha_i\mathbf{B}\right] \;\forall\; i$, he transforms the concentrated model in (6) into

$$R_{0,i} - \alpha_i I_r R_{1,i}^{(1)} = \alpha_i \mathbf{B} R_{1,i}^{(2)} + U_i \quad \text{with} \quad R_{1,i} = \begin{bmatrix} R_{1,i}^{(1)} \\ R_{1,i}^{(2)} \end{bmatrix} \tag{23}$$

[11] See Vlaar (2004), who imposes linear short- and long-run restrictions with restriction matrices for the cases of just- and over-identification. Compare also to Hamilton (1994, Ch. 11, p. 331), Lütkepohl (2006, Ch. 3.2, p. 84), and Kilian and Lütkepohl (2017, Ch. 11.2.2 and p. 315), who present this as full-information MLE.

[12] See also Lütkepohl (2005, Ch. 7.2.2) or Brüggemann and Lütkepohl (2005) about GLS estimation of the individual cointegration parameters. In fact, (25) reduces to this GLS estimator if there is only $N = 1$ individual entity. Moreover, Lütkepohl (2005, Ch. 7.3.2) and Breitung (2005, p. 159) consider restricted β.

partitioned into $R_{1,i}^{(1)}$ as the r upper time series and $R_{1,i}^{(2)}$ as the remaining $r_\perp$ time series and n_{d1} restricted deterministic regressors. The left-multiplication $\alpha_i^{(+)}\alpha_i = I_r$ eliminates the individual loadings α_i in (23) such that (6) is further transformed into

$$\alpha_i^{(+)}\left(R_{0,i} - \alpha_i R_{1,i}^{(1)}\right) = \mathbf{B}R_{1,i}^{(2)} + \alpha_i^{(+)}U_i \quad \text{using} \quad \alpha_i^{(+)} = \left(\alpha_i^\top \Sigma_{u,i}^{-1}\alpha_i\right)^{-1}\alpha_i^\top\Sigma_{u,i}^{-1}. \tag{24}$$

The *pooled OLS estimator* of the homogeneous $\mathbf{B}$, i.e. the right $r \times (r_\perp + n_{d1})$ block of the normalized and transposed cointegrating matrix, yields the second step given by

$$\widehat{\mathbf{B}}_{2S} = R_0^{(+)}R_1^{(2)\top}\left(R_1^{(2)}R_1^{(2)\top}\right)^{-1} \tag{25}$$

$$\begin{aligned} &\text{with regressand} \quad R_{0,i}^{(+)} := \alpha_i^{(+)}R_{0,i} - R_{1,i}^{(1)} \quad &&\text{collected in} \quad R_0^{(+)} := \left[R_{0,1}^{(+)} : \cdots : R_{0,N}^{(+)}\right] \\ &\text{and regressor} \quad R_{1,i}^{(2)} &&\text{collected in} \quad R_1^{(2)} := \left[R_{1,1}^{(2)} : \cdots : R_{1,N}^{(2)}\right]. \end{aligned}$$

For feasible estimation, inserting any $\sqrt{T}$-consistent estimates for α_i and $\Sigma_{u,i}$ leaves the limiting distribution of the estimator in (25) unaffected. This first step may be the individual ML estimation of each VECM, which concentrates out the short-run effects and unrestricted deterministic terms $\Gamma_i Z_{2,i}$ from the observed time series as well. Under rank restriction r, the second-step estimates $\widehat{\mathbf{B}}_{2S}$ are thereby $T\sqrt{N}$-consistent (Breitung 2005, p. 156, Th. 1). Overall, this procedure thus follows the steps of the ML estimation, but sets a pooled estimation on top of the individual reduced-rank regressions of (8). Like in third step of the individual ML estimation, the cointegrating matrix $\widehat{\beta}^\top = \left[I_r : \widehat{\mathbf{B}}_{2S}\right]$ can be reintroduced into the conditional OLS of (9) of the individual parameters, for which product moments are already available from the first step. Thereof, the estimates for α_i and $\Sigma_{u,i}$ – now conditional on the pooled $\widehat{\beta}$ – can be used in (25) again. Hlouskova and Wagner (2010, p. 195) expand this principle and iterate conditional OLS and second-step estimation alternately until reaching convergence of the estimates $\widehat{\mathbf{B}}_{2S}$.

Deterministic term. Since the first estimation step by (6) removes the unrestricted regressors $\boldsymbol{d}_{2,it}$ from the VECM in (2), the coefficients $\Phi_{2,i}$ are individual-specific by construction. In consequence, the homogeneity restriction $\beta_{K,i} = \beta_K$ in

$$\Pi_i := [\Pi_{K,i} : \Phi_{1,i}] = \alpha_i\left[\beta_K^\top : \beta_{0i}^\top\right] = \alpha_i\left[I_r : \mathbf{B}^{(2)} : \mathbf{B}_i^{(3)}\right] \tag{26}$$

implies one additional choice for specifying the innovative deterministic term in Table 2:[13] The coefficients for $D_{1,i}M_{2,i}$, containing the restricted deterministic regressors $\boldsymbol{d}_{1,it}$, can be defined as having some homogeneous $\beta_{0i} = \beta_0\ \forall\ i$ or individual-specific effects β_{0i}. For example, *Case 2* with heterogeneous intercepts β_{0i} and $p_i = 1 \forall i$, i.e. without the need to correct for lagged short-run effects by $M_{2,i}$, adds *fixed effects* to the auxiliary model of (24). The regressors of such individual-specific $\mathbf{B}_i^{(3)}$ are partialled out from the pooled regression beforehand, while $\mathbf{B}^{(2)}$ is simply extended by the $r \times n_{d1}$ homogeneous $\beta_0^\top$ and estimated by the pooled $\widehat{\mathbf{B}}_{2S}$ of (25). Note however that $\Phi_{1,i} = \alpha_i\beta_0^\top$ remains heterogeneous irrespective of the homogeneity restriction on β_0.

[13] Groen and Kleibergen (2003, Ch. 4.1) give an overview of the different combinations that emerge from this single additional option.

Testing the cointegration rank

Like the individual tests in Section 2.1.2, the following panel tests evaluate the hypothetical cointegration rank $r_{H0} = 0, \ldots, K-1$ within a sequential testing procedure. Each of the N individuals must have observations on the same K variables so that the hypothesis pair on the heterogeneous ranks r_i in the individual VAR processes $\boldsymbol{y}_{it}$ can be stated as

$$\begin{aligned} H_0 : \bar{r} = r_{H0} \quad \text{versus} \quad H_1 : \operatorname{rk}(\Pi_i) > r_{H0} \text{ for some } i \\ \text{with } \bar{r} = \max\{\operatorname{rk}(\Pi_i) \mid i = 1, \ldots, N\}. \end{aligned} \tag{27}$$

While a statistical heterogeneity in r_i lets the sequential procedure find the maximum rank among the individual processes, the theoretical context might allow for the more restrictive assumption of homogeneous cointegration ranks across all individuals $i = 1, \ldots, N$. Accordingly, the hypothesis pair in (27) would adopt a common rank $r_i = \bar{r} = r$. Carrion-i Silvestre and Surdeanu (2011, p. 9) assume this "for the panel data based procedure to make sense" and the homogeneous cointegrating vectors $\beta_{K,i} = \beta_K$ in Breitung (2005) and Groen and Kleibergen (2003) actually have the statistical implication of $r_i = r$.

Pooled cointegrating vectors. The more restrictive assumption of homogeneous β_K allows for a pooled estimation to further increase the test power. Only in this case, the *pooling* and the *combination* approach of the panel test differ. After estimating individual models with pooled β_K, Breitung (2005) combines the individual LM-test results of (17) by their averaged statistics in (28). Originally, he does not make use of the gamma distribution to derive p-values of the LM-tests. Since this option follows straightforwardly from the asymptotic equivalence of the LM- and LR-test, **pvars** implements their meta-analytical combination, too.[14] Thereby, both combination approaches are available to shrink the vectors of N test results into scalar decisions. The following section refers to these combination approaches, which are congruent with the pooling approaches of Table 1.

Combination approaches. Extending the combination idea from Im *et al.* (2003) to the multivariate case of K variables, Larsson *et al.* (2001) use the *cross-sectional average* of individual test statistics $\lambda_i^\bullet$, which are standardized by the first and second moment. Accordingly, the so-called LR-bar test statistic $\Upsilon_{\overline{LR}}$ is calculated as

$$\begin{aligned} \overline{\lambda}(r_{H0}) &= \frac{\sum_{i=1}^{N} \lambda_i^\bullet(r_{H0})}{N}, \\ \Upsilon_{\overline{LR}} &= \frac{\sqrt{N}\left(\overline{\lambda}(r_{H0}) - \mathbb{E}\left(Z_{r_\perp}\right)\right)}{\sqrt{\operatorname{Var}\left(Z_{r_\perp}\right)}} \xrightarrow{d} \mathcal{N}(0,1) \end{aligned} \tag{28}$$

and follows asymptotically a standard normal distribution under H_0 as $T \to \infty$ and subsequently $N \to \infty$. This holds also if $T, N \to \infty$ simultaneously, but T has a faster limiting rate such that $\sqrt{N}/T \to 0$. The combination approach itself is well-established for all kinds of panel tests. The key contribution by Larsson *et al.* (2001) is to prove that the moments of the asymptotic distribution $Z_{r_\perp}$ exist.[15] Likewise, Örsal and Droge (2014) present and utilize a proof for the detrending panel SL-test with the deterministic type *SL_trend*. Additionally,

[14] Choi (2001, p. 268) shows for the univariate case of panel unit root tests that especially the inverse normal test has a favorable power in comparison to the panel test of averaged statistics.

[15] See also the corrigendum by Örsal and Droge (2011).

pvars offers the specification *SL_mean* because the existence of the related moments follows directly from the identity of $Z_{r_\perp}$ for *SL_mean* and *Case 1* – see (19).

Following the *meta-analytical combination* of p-values, Choi (2001) uses different combination methods on the individual p-values denoted by p_i. The first combination test, which Fisher (1932) originally proposed and Maddala and Wu (1999) embraced for tests in nonstationary panels, is the inverse chi-square test. Its statistic is stated as

$$P = -2\sum_{i=1}^{N} \ln\left(p_i\right) \xrightarrow{d} \chi^2(2N) \tag{29}$$

under the asymptotics of $T_i \to \infty$ but fixed N. In the limit of N, however, the distribution of P degenerates. For panels with many[16] individuals, Choi (2001) thus standardizes P by the moments of the χ^2-distribution and proposes the modified statistic

$$\begin{aligned} P_m &= -\frac{\sum_{i=1}^{N}\left(\ln\left(p_i\right)+1\right)}{\sqrt{N}} \\ &= \frac{P-2N}{\sqrt{4N}} \xrightarrow{d} \mathcal{N}(0,1), \end{aligned} \tag{30}$$

when $T_i \to \infty$ and then additionally $N \to \infty$. Furthermore, assuming the same limiting data dimensions, Choi (2001) uses the inverse normal test

$$Z = \frac{\sum_{i=1}^{N}\Phi_{\mathcal{N}}^{-1}\left(p_i\right)}{\sqrt{N}} \xrightarrow{d} \mathcal{N}(0,1) \tag{31}$$

from Stouffer, Suchman, DeVinney, Star, and Williams (1949) for nonstationary panel data analysis. Here, $\Phi_{\mathcal{N}}^{-1}(\bullet)$ denotes the probit function, i.e. the inverse of the standard normal cumulative distribution.

PANIC. The *Panel Analysis of Nonstationarity in Idiosyncratic and Common components* by Bai and Ng (2004) accounts for cross-sectional dependence stemming from unobserved common factors, which could be stationary, integrated, or both coexistent. Arsova and Örsal (2017; 2018) employ PANIC on VAR systems so that their multivariate extension then allows for tests on the cointegration rank within the idiosyncratic VECM of the individuals. The additive model is stated as

$$\begin{aligned} \boldsymbol{y}_{it} &= \Lambda_i^\top \boldsymbol{F}_t + \boldsymbol{y}_{it}^{id}, \\ \boldsymbol{y}_{it}^{id} &= \boldsymbol{\mu}_{0i} + \boldsymbol{\mu}_{1i} t + \boldsymbol{y}_{it}^{dt}, \\ \boldsymbol{y}_{it}^{dt} &= A_{i1}\boldsymbol{y}_{i,t-1}^{dt} + \ldots + A_{i,p_i}\boldsymbol{y}_{i,t-p_i}^{dt} + \boldsymbol{u}_{it}^{dt} \\ (1-L)\boldsymbol{F}_t &= C(L)\boldsymbol{u}_t^F. \end{aligned} \tag{32}$$

The idea of PANIC is to perform separate analyses on the common factor $\boldsymbol{F}_t$ and the idiosyncratic component $\boldsymbol{y}_{it}^{id}$, both being part of the observable multivariate time series $\boldsymbol{y}_{it}$. Hence, factors $\boldsymbol{f}_t$ and heterogeneous loadings Λ_i are first estimated via PCA on the first-differenced data matrix of dimension $(T-1) \times (K \cdot N)$. If unknown, the number of underlying factors

[16] Choi (2001, p. 268) states for panel unit root tests that, even with samples of $N = 100$, P_m does not follow its asymptotic distribution closely. In contrast to the two Fisher tests, the test size of the inverse normal test and of the averaged statistics is more robust to any choice of N.

can be chosen according to information criteria by Bai and Ng (2002) or Onatski (2010) as illustrated in Section 4.1. The factors $\hat{\boldsymbol{F}}_t = \sum_{j=1}^{t} \hat{\boldsymbol{f}}_j$ with initial $\hat{\boldsymbol{f}}_1 = \boldsymbol{0}_L$ are re-accumulated into levels to estimate the idiosyncratic component $\hat{\boldsymbol{y}}_{it}^{id} = \boldsymbol{y}_{it} - \hat{\Lambda}_i^\top \hat{\boldsymbol{F}}_t$. While PCA generates orthogonal factors $\hat{\boldsymbol{f}}_t$ normalized to an identity covariance matrix, the re-accumulated factors $\hat{\boldsymbol{F}}_t$ are usually *oblique* and described by a single VAR model. The VAR process of $\boldsymbol{F}_t$ can further contain common structural breaks from $\boldsymbol{y}_{it}$ such that, for instance, Örsal (2017) resorts to JMN (2000) for testing the cointegration rank of $\hat{\boldsymbol{F}}_t$.

Since $\boldsymbol{F}_t$ is assumed to be the exclusive source of cross-sectional dependence, the idiosyncratic VECM of $\boldsymbol{y}_{it}^{id}$ are independent across individuals i and the basic methodology of first-generation tests is valid again. Örsal and Arsova (2017) use the approach of meta-analytical combination of the idiosyncratic p-values and Arsova and Örsal (2018) the averaged test statistics. Accordingly, the nested hypotheses of (27) about r_{H0} refer to idiosyncratic cointegration within $\boldsymbol{y}_{it}^{id}$ and, for conclusions on cointegration within the observable $\boldsymbol{y}_{it}$, the nonstationary properties of $\boldsymbol{F}_t$ need to receive attention. Individual unit root and cointegration tests can be applied to the common factors $\boldsymbol{F}_t$ and their VAR process. If all series in $\boldsymbol{F}_t$ are stationary, panel test results on $\boldsymbol{y}_{it}^{id}$ hold equally for $\boldsymbol{y}_{it}$. If $\boldsymbol{F}_t$ contains stochastic trends, those enter the observable variables $\boldsymbol{y}_{it}$, too. In this case, they can confound the interpretation, for example, if some of the heterogeneous loadings Λ_i are zero and thus do not let stochastic trends in $\boldsymbol{F}_t$ drive the variables $\boldsymbol{y}_{it}$ equally.

For testing the cointegration rank in each idiosyncratic component, Arsova and Örsal (2017; 2018) consider two individual procedures: The Johansen test applied to $\boldsymbol{y}_{it}^{id}$ directly and the SL-test applied to $\boldsymbol{y}_{it}^{dt}$ after the GLS-based trend adjustment. The SL-test based on the defactored and detrended series $\boldsymbol{y}_{it}^{dt}$ assesses only the null hypothesis of no cointegration. For $r_{H0} > 0$ in the sequential test, the $r_\perp = K - r_{H0}$ stochastic trends in the rank-restricted VECM are calculated as $(\beta_{i\perp})^\top \boldsymbol{y}_{it}^{id}$. Arsova and Örsal (2017; 2018) then apply the SL-test to the presumably $r_\perp$ stochastic trends under the H_0 of no cointegration in order to test the equivalent H_0 of rank r_{H0} in the idiosyncratic process of $\boldsymbol{y}_{it}^{id}$.

Defactoring adds a linear trend to the estimated series $\hat{\boldsymbol{y}}_{it}^{id}$ inevitably, which affects the distribution of $Z_{r_\perp}$ for both, the SL- and the Johansen test. For deriving individual p-values via the gamma approximation as in (20) or for standardizing the LR-bar panel test statistic as in (28), the panel tests rely on the moments of $Z_{r_\perp}$ which Arsova and Örsal (2018, Appendix, Tab. A.1) have simulated and tabulated. The moments depend on $r_\perp$ and the prescribed deterministic trend, but not on the common factors $\boldsymbol{F}_t$. Likewise, they can be calculated via the response surface regression model by Trenkler (2008) using the coefficients for the case of a deterministic trend.[17] Note that Örsal and Arsova (2017) did not originally propose combining individual p-values from defactored Johansen tests. However, **pvars** implements this in accordance with the convergence result of $Z_{r_\perp}$ implied by Arsova and Örsal (2018, p. 1041, Th. 3.4).

Correlated probits. Cross-sectional dependence between the individual entities induces correlation between the probits $\Phi_{\mathcal{N}}^{-1}(p_i)$ of the individual cointegration tests. In order to correct for this and robustify the inverse normal test of (31), Arsova and Örsal (2021) combine individual p-values from the SL-procedure with Hartung's (1999) modifications and with

[17] For an illustration, compare the results of the R commands:

```
R> pvars:::coint_moments$SL_trend[12:1, ]
R> pvars:::aux_CointMoments(dim_K=12, rs_coef=pvars:::coint_rscoef[["SL_trend"]])
```

their own *correlation-augmented inverse normal* (CAIN) test. The panel test statistic for the modified combination approach is given by

$$\begin{aligned} t(\tilde{\rho}) &= \frac{\sum_{i=1}^{N} \Phi_{\mathcal{N}}^{-1}(p_i)}{\sqrt{N + (N^2 - N) \cdot \tilde{\rho}^{\bullet}_{\text{probit}}}} \xrightarrow{d} \mathcal{N}(0,1) \quad \text{with} \\ \tilde{\rho}^{\text{HA}}_{\text{probit}} &= \hat{\rho}^{*}_{\text{probit}} + \kappa \cdot \sqrt{\frac{2}{(N+1)}} \left(1 - \hat{\rho}^{*}_{\text{probit}}\right) \quad \text{or} \\ \tilde{\rho}^{\text{CAIN}}_{\text{probit}} &= g\left(\rho_{\epsilon}, K, r_{H0}\right). \end{aligned} \tag{33}$$

The correction factor $\tilde{\rho}^{\text{HA}}_{\text{probit}}$ by Hartung (1999) is based on his unbiased and consistent estimator $\hat{\rho}^{*}_{\text{probit}} = \max\{-\frac{1}{N-1},\ 1 - s^2_{N-1}\left(\Phi_{\mathcal{N}}^{-1}(p_i)\right)\}$ of the probit correlation, where $s^2_{N-1}(\cdot)$ denotes the unbiased sample variance. For the scaling factor κ, he proposes $\kappa_1 = 0.2$ and, additionally, $\kappa_2 = 0.1 \cdot \left(1 + \frac{1}{N-1} - \hat{\rho}^{*}_{\text{probit}}\right)$ "working mainly for smaller $\hat{\rho}^{*}_{\text{probit}}$" (Hartung 1999, p. 853). He does not state any decision criterion to distinguish between the intervals of smaller and larger values of $\hat{\rho}^{*}_{\text{probit}}$ however.

The correction factor $\tilde{\rho}^{\text{CAIN}}_{\text{probit}}$ by Arsova and Örsal (2021) is the empirical estimate of the average correlation between the individual probits. For translating cross-sectional dependence within the data panel into $\tilde{\rho}^{\text{CAIN}}_{\text{probit}}$, the authors provide response surface regression coefficients for the regression model $g\left(\rho_{\epsilon}, K, r_{H0}\right)$. Therein, the number of endogenous variables K and the tested cointegration rank r_{H0} follow directly from the model resp. test specification. The *average absolute pairwise cross-sectional correlation* ρ_{ϵ} between the residuals is estimated for $r_{H0} = 0$ only and then used for all r_{H0} of the sequential testing procedure up to $r_{H0} = K - 1$. The estimator for this mean sample correlation of the residuals within the *same* variable $k = 1, \ldots, K$ and between the different individuals is

$$\hat{\rho}_{\epsilon} = \frac{2}{K \cdot N \cdot (N-1)} \sum_{k=1}^{K} \sum_{i=1}^{N} \sum_{j=i+1}^{N} |\hat{\rho}_{ki,kj}|. \tag{34}$$

Since Arsova and Örsal (2021) assume that the pairwise correlations between the residuals of *different* variables converge to zero for $N \to \infty$, the sample correlations $\hat{\rho}_{ki,k^*j}$, $k \neq k^*$, are omitted in (34). In doing so, the cross-sectional correlation between variables does not reduce the average of the presumably stronger "within"-correlations, which mitigates the hazard of underestimating ρ_{ϵ}. Only if the empirical "between"-correlation surpasses "within"-correlation, $\hat{\rho}_{\epsilon}$ could actually understate cross-sectional dependence.

If the individual, potentially heterogeneous break periods $\boldsymbol{\tau}_i$ and the number of breaks are known, this third-generation panel test can also respect trend breaks in the cointegration relationship. For this, the individual GLS-based trend-adjustment by Trenkler *et al.* (2008) removes the deterministic component from the observed time series of (10) including up to two trend breaks. Then, the standard LR-test procedure is applied under consideration of the trend-break specific test distribution. The individual p-values are combined as described in (33) in order to account for the cross-sectional dependence.

Identifying structure


In comparison to the individual time series analysis, the cross-section dimension of the panel data enables additional ways to identify structural shocks $\boldsymbol{\epsilon}_{it} = \mathsf{B}_i^{-1}\boldsymbol{u}_{it}$ from the K reduced-form errors $\boldsymbol{u}_{it}$ of the VAR model in (1). The identification across the individuals $i = 1, \ldots, N$

may start with normalizing $\boldsymbol{u}_{it}$ to unit variance by $\mathsf{B}_{\sigma,i} = \text{diag}\,(\sigma_{u,1i}, \ldots, \sigma_{u,Ki})$ such that the structural decomposition concentrates on the residual correlation matrix

$$\begin{aligned} \Omega_{u,i} &:= \mathsf{B}_{\sigma,i}^{-1} \Sigma_{u,i} \mathsf{B}_{\sigma,i}^{-1} = \mathsf{B}_{\Omega,i} \mathsf{Q}_i \mathsf{Q}_i^\top \mathsf{B}_{\Omega,i}^\top \\ \text{with } \mathsf{B}_{\Omega,i} &:= \text{chol}\,(\Omega_{u,i}) \text{ and } \mathsf{Q}_i \mathsf{Q}_i^\top = \mathsf{Q}_i \mathsf{Q}_i^{-1} = I_K. \end{aligned} \tag{35}$$

The usual Cholesky decomposition $\mathsf{B}_i\,(\boldsymbol{0}) := \text{chol}\,(\Sigma_{u,i}) = \mathsf{B}_{\sigma,i} \cdot \text{chol}\,(\Omega_{u,i})$ generates *whitened* shocks $\boldsymbol{\epsilon}_{\boldsymbol{0}it} = \mathsf{B}_i\,(\boldsymbol{0})^{-1} \boldsymbol{u}_{it}$, which already conform with the defining $\boldsymbol{\epsilon}_{\boldsymbol{0}it} \sim (0, I_K)$ after scaling $\left(\mathsf{B}_{\sigma,i}^{-1}\right)$ and de-correlating $\left(\mathsf{B}_{\Omega,i}^{-1}\right)$ the zero-mean vector $\boldsymbol{u}_{it}$. Further, matrix Q_i is orthogonal and can be a product of $K(K-1)/2$ Givens rotation matrices, which map from the tuple of rotation angles $\boldsymbol{\theta}_i$ to the total set of candidate shocks $\boldsymbol{\epsilon}_{\boldsymbol{\theta}it} = \mathsf{Q}\,(\boldsymbol{\theta}_i)^{-1} \boldsymbol{\epsilon}_{\boldsymbol{0}it}$ and their impact matrices $\mathsf{B}_i\,(\boldsymbol{\theta}_i) = \mathsf{B}_i\,(\boldsymbol{0})\,\mathsf{Q}\,(\boldsymbol{\theta}_i)$. As the identification procedures rely on estimated $\widehat{\boldsymbol{u}}_{it}$, the candidate samples $\mathcal{A}_{\boldsymbol{\theta}}^{(\mathrm{i})} = \{\widehat{\boldsymbol{\epsilon}}_{\boldsymbol{\theta}it}\}_{t=1}^{T}$ of dimension $K \times T$ involve the usual issues of individual SVAR with small or medium-sized time series. The panel methods presented in the following offer the advantage of *(i)* increasing the power of dependence tests in ICA and of *(ii)* accommodating structural information from cross-sectional dependence.

Figure 1: Panel SVAR identification by ICA in Herwartz and Wang (2024).



Common rotation. Herwartz and Wang (2024) propose and evaluate the pooled identifica-

tion procedure stylized in Figure 1. An individual decomposition $\boldsymbol{\epsilon}_{\boldsymbol{\theta}it} = \mathsf{Q}(\boldsymbol{\theta})^{-1} \mathsf{B}_{\Omega,i}^{-1} \mathsf{B}_{\sigma,i}^{-1} \boldsymbol{u}_{it}$ allows for a common rotation $\mathsf{Q}(\boldsymbol{\theta})$ of the whitened shocks while accounting for heterogeneous variances and cross-variable correlations in $\boldsymbol{u}_{it}$. Among competing *pooled samples* $\mathcal{A}_{\boldsymbol{\theta}}^{(N)} = \{\{\widehat{\boldsymbol{\epsilon}}_{\boldsymbol{\theta}it}\}_{t=1}^{T}\}_{i=1}^{N}$ of dimension $K \times (T \cdot N)$, the ICA then determines the least dependent shocks $\widehat{\boldsymbol{\epsilon}}_{\widehat{\boldsymbol{\theta}}it}$ with optimal $\widehat{\boldsymbol{\theta}}$, from which $\widehat{\mathsf{B}}_i$ is recovered for each individual.

Common shocks. Calhoun *et al.* (2001) propose *group ICA* and apply this to panel data originating from functional Magnetic Resonance Imaging (fMRI) of brains. In the same empirical context, Risk, Matteson, Ruppert, Eloyan, and Caffo (2014) evaluate ICA algorithms, which can be applied to the panel of reduced-form errors $\boldsymbol{u}_{it}$ alike. Accordingly, their model $\boldsymbol{u}_{it} = \mathsf{B}_i \boldsymbol{\epsilon}_t + \boldsymbol{e}_{it}$ consists of L common shocks $\boldsymbol{\epsilon}_t$ and some idiosyncratic noise $\boldsymbol{e}_{it}$ (Risk *et al.* 2014, p. 227). In a two-step PCA, they first whiten each individual sample $[\widehat{\boldsymbol{u}}_{i1} : \ldots : \widehat{\boldsymbol{u}}_{iT}]^\top = \mathsf{U}_i \mathsf{D}_i \mathsf{V}_i^\top$ by compact *singular value decomposition* such that $\mathcal{A}_i := \sqrt{T}\mathsf{U}_i^\top$ has an identity covariance matrix I_K. They further factorize the $T \times (K \cdot N)$ concatenated samples $\left[\mathcal{A}_1^\top : \ldots : \mathcal{A}_N^\top\right] = \mathsf{U}\mathsf{D}\mathsf{V}^\top$ and utilize the first L columns of left singular vectors to construct an $L \times T$ baseline sample $\mathcal{A}_{\mathbf{0}}^{(2S)} := \sqrt{T}\mathsf{U}_{1:L}^\top$. The ICA, which becomes noise-free after this data reduction, then determines the least dependent common shocks $\widehat{\boldsymbol{\epsilon}}_{\widehat{\boldsymbol{\theta}}t}$. The multivariate least squares regression of $\widehat{\boldsymbol{u}}_{it}$ on $\widehat{\boldsymbol{\epsilon}}_{\widehat{\boldsymbol{\theta}}t}$ recovers the $K \times L$ impact matrices $\widehat{\mathsf{B}}_i$, $\forall i = 1, \ldots, N$.

ICA. If at most one of the shocks is Gaussian, the *independent component analysis* as established by Comon (1994) can determine the rotation angles $\widehat{\boldsymbol{\theta}}^{\bullet}$ and thereby identify the impact matrix $\widehat{\mathsf{B}}_i^{\bullet}$ of the methods $\bullet \in \{(i), (N), (2S)\}$. For this purpose, dependence measures $\mathcal{D}(\cdot)$ discriminate between the candidates of whitened shocks $\mathcal{A}_{\boldsymbol{\theta}}^{\bullet} = \mathsf{Q}(\boldsymbol{\theta})^{-1} \mathcal{A}_{\mathbf{0}}^{\bullet}$ by dependencies beyond the second moment. In the spirit of Hodges-Lehmann estimation (1963), minimization procedures find the rotation angles $\widehat{\boldsymbol{\theta}}^{\bullet} = \operatorname{argmin}_{\boldsymbol{\theta}} \mathcal{D}(\mathcal{A}_{\boldsymbol{\theta}}^{\bullet})$ of the least dependent shocks.

ICA-based identification procedures for the individual SVAR are already implemented in **svars**. As we focus on their embedding into a panel framework, we just list them here briefly and refer the reader to the accompanying vignette (Lange *et al.* 2021) for a comprehensive overview and to the Monte Carlo study (Herwartz, Lange, and Maxand 2022) for a performance assessment. The following dependence measures $\mathcal{D}(\cdot)$ and optimization procedures are adopted in **pvars**: The *Cramér-von Mises* (CvM) *distance* by Genest *et al.* (2007) is used in **svars**' two-step optimization procedure with **copula** (Kojadinovic and Yan 2010) and has been exemplarily applied for individual SVAR by Herwartz (2018). The *distance covariance* (dCov) by Székely *et al.* (2007) is used in the gradient algorithm of **steadyICA** (Risk, James, and Matteson 2015) and has been applied for SVAR by Matteson and Tsay (2017). The *dependence coefficient* (dCoef) by Bakirov *et al.* (2006) is not used in **svars** and **pvars**.[18]

ICA can identify shocks and impact matrices up to scaling, column signs, and ordering only. For example in the case of $K = 2$, the relevant interval of a full rotation $\theta \in (0, 2\pi]$ reduces to a quadrant, e.g. $\theta \in (0, \pi/2]$, since any exceeding rotation just permutes the ordering and reverses signs. If $\theta \in (\pi/2, 2\pi]$, the results of $\mathcal{D}(\mathcal{A}_{\boldsymbol{\theta}}^{\bullet})$ including the minima would be identical to those of the first quadrant. Against this ambiguity, a common practice for the unique identification of B_i is to *(1)* choose the column ordering which maximizes the sum of the absolute diagonal elements and *(2)* then switch signs of those columns whose main diagonal element turns out to be negative. Under $\boldsymbol{\epsilon}_{it} \sim (0, I_K)$, each shock is thereby attributed to the variable on which it has the strongest effect on impact.

[18] Note that dCoef and dCov are implemented in **energy** by Rizzo and Szekely (2022).

3. Implementation

For each field of VAR application and for the supporting tools, Table 3 displays **pvars**' core functions, the S3-class of their output object and their dependencies within and to other packages. In particular, several classes and their corresponding methods are imported from **svars**. In addition to the familiar methods such as `print()` and `summary()`, **pvars** offers the method `toLatex()` for conveniently formatting **pvars** results into LaTeX objects, thus minimizing the risk of reporting errors from tedious copy-pasting.

Table 3: Package design of **pvars**.


Function	Class	Branch	Methods	Description	Literature
3.1 Testing the cointegration rank					
`pcoint.JO`	`pcoint`	`coint.JO`	`print`,	Panel Johansen tests	Larsson *et al.* (2001), Choi (2001)
`pcoint.BR`	`pcoint`	`coint.JO`	`summary`,	Panel test with pooled β	Breitung (2005)
`pcoint.SL`	`pcoint`	`coint.SL`	`toLatex`	Panel SL-tests	Örsal, Droge (2014)
`pcoint.CAIN`	`pcoint`	`coint.SL`		Correlation-augmented tests	Arsova, Örsal (2021), Hartung (1999)
3.2 Estimating VAR models					
`pvarx.VAR`	`pvarx`	`VAR`[a)]	`irf`, `print`,	Mean-group estimation	Rebucci (2010), Pesaran, Smith (1995)
`pvarx.VEC`	`pvarx`	`VECM`	`summary`	Pooled cointegrating vectors β	Breitung (2005), Pesaran *et al.* (1999)
3.3 Identifying structure					
`pid.chol`	`pid`	`id.chol`[a)]	`irf`,	Recursive causality	Sims (1980)
`pid.grt`	`pid`	`id.grt`[c)]	`print`,	Long- & short-run restrictions	Breitung *et al.* (2004)
`pid.iv`	`pid`	`id.iv`	`summary`,	Proxy SVAR	Empting et al. (2026)
`pid.dc`	`pid`	`id.dc`[a)]	`toLatex`	ICA by distance covariance	Calhoun *et al.* (2001) and
`pid.cvm`	`pid`	`id.cvm`[a)]		ICA by Cramér-von Mises dist.	Herwartz, Wang (2024)
3.4 Supporting tools					
`speci.factors`	`speci`	—	`print`	Criteria for the number of factors	Bai, Ng (2002; 2004), Onatski (2010)
`speci.VAR`	`speci`	—	`print`	Criteria for lags p and periods τ	Bai, Perron (1998; 2003), Yang (2002)
`irf.pvarx`	`svarirf`[b)]	`irf.varx`	`plot`, `print`	Mean-group IRF [S3-METHOD]	Sims (1980), Gambacorta et al. (2014)
`PP.system`	`svarirf`[b)]	—	`plot`, `print`	Persistence profiles	Lee, Pesaran (1993) and
`PP.variable`	`svarirf`[b)]	—	`plot`, `print`	(Structural) persistence profiles	Pesaran, Shin (1996)
`sboot.mg`	`sboot`[b)]	—	`print`, `summary`,	Mean-group inference	Pesaran, Smith (1995)
`sboot.pmb`	`sboot`[b)]	`sboot.mb`	`plot`, `toLatex`	Panel-block bootstrap	Empting *et al.* (2026)

[a)] Consider the R packages **vars** and **svars** for these functions. Like `id.dc`, **pvars**' `pid.dc` uses **steadyICA** by Risk *et al.* (2015).
[b)] This class and its methods are imported from the R package **svars** by Lange *et al.* (2021).
[c)] Its scoring algorithm is part of the `SVEC` function from the R package **vars** by Pfaff (2008b).

Figure 2 illustrates **pvars**' modular implementation, which leads to the three layers of dependency between the library of auxiliary functions, the functions for individual econometric procedures, and their panel extensions. In this tree-like structure, the `aux`-functions are the hidden "roots" and the individual functions are "leafy branches". Some branches of the individual `id.*()` functions are a "graftage" from **vars** and **svars**. The individual `coint.*()` functions are more flexible than existing implementations of cointegration rank tests and extend the set of available specification options. Finally attached to the branches, the "fruits" of this package are the panel functions for each field of VAR application. These are printed in bold to emphasize their novelty and **pvars**' main contribution.

The idea of modular programming is to break monolithic and repetitive code down into functional sub-entities, which achieves easier maintenance, better testability, and reusability. Especially the multivariate panel procedures benefit from this kind of implementation because their functions reduce to simple re-arrangements of auxiliary functions. These can be repeatedly applied over subsets of the $K \times T \times N$ data array. Whenever sensible, the R functions are vectorized in order to enable more flexible input arguments and faster matrix computation of the multivariate VAR models. With automatic unit tests by **testthat** (Wickham 2011), `pcoint.*()` and subordinated modules are checked against reproduced examples of the literature. Functions outside this hierarchy are checked against consistency with the

Figure 2: Modular implementation in three layers.



other **vars** packages, identities derived from econometric theory, and simulation results from Empting *et al.* (2026).

Argument structure. In order to comply with the modular functions and their repeated calls over data subsets, several conventions have been established for the input arguments of the auxiliary, individual, and panel functions. The arguments whose names are marked with the prefix `L.*` are an R object of class `list` with N elements of repeated structure. Most prominently, `L.data` is the panel data in *list format* and contains N `data.frame`s of dimension $T_i \times K$. In these multivariate time series, the variables must have an identical order $k = 1, \dots, K$ for each individual $i = 1, \dots, N$. All successions are binding for any other `L.*` object within the environment of a given function. Hence even in the global environment, the user should stick to a universal succession for all `L.*` objects in order to avoid confusion.

Further conventions for prefixes are `n.*` and `dim_*`, which denote integers and refer to a quantity resp. the dimension of a matrix or array. Straightforwardly, `dim_K`, `dim_T`, and `dim_N` signify the data dimensions K, T, and N for instance. Arguments with prefix `t_*` define $\boldsymbol{\tau}$ for the period-specific deterministic regressors. As a `list`, they collect optional vectors of integers for trend breaks (named `t_break`), shift dummies (`t_shift`), and impulse dummies (`t_impulse`) as well as a single integer `n.season`, which indicates the seasonal frequency for the centered dummies. Each integer in those vectors specifies a period τ at which a break, shift, or impulse occurs within the interval $1, \dots, T$ of the complete panel data set (i.e. including the presamples). The following sections describe how the mandatory and optional input arguments are employed in the **pvars** functions for cointegration tests (Section 3.1), estimation (Section 3.2), and structural identification (Section 3.3).

3.1. Testing the cointegration rank

The panel cointegration tests are implemented in accordance with the three dimensions of panel test construction as presented in Table 1. *(i)* Each `pcoint.*()` function always assesses all hypotheses $r_{H0} = 0, \dots, K-1$ and uses all combination approaches available for the respective underlying individual procedure. *(ii)* The data generating process by contrast, which covers also the panel test generation, must be defined via the arguments of `pcoint.*()`. These arguments are kept consistent across the functions or refer to the underlying procedure. *(iii)* The panel functions `pcoint.*()` usually loop over their internal function `cointf()` as the underlying procedure. `cointf()` arranges the auxiliary functions in the same way as its respective branch `coint.*()`, but has been modified for the panel test. Hence, if an individual sample does not conform with the panel result, the user may consult the individual counterpart

`coint.*()` and the persistence profiles `PP.*()` for a closer inspection of the peculiar individual entity. Within the scope of PANIC, the test functions `coint.*()` for the single VAR system can assess cointegration between $\boldsymbol{F}_t$ from (32), too.

The branch of Johansen. The individual Johansen procedure is available as the function

```
coint.JO(y, dim_p, x, dim_q, type, t_D1=NULL, t_D2=NULL)
```

By default, the R function performs both LR-test variants on the multivariate time series `y` with lag order `dim_p`. For the *trace test* as in (15) and exclusively for the individual *maximum eigenvalue test* with simple specifications, p-values are approximated by the gamma distribution. Optionally, weakly exogenous variables `x` with lag order `dim_q` are incorporated. The character argument `type` defines the conventional deterministic term according to the labels for the *innovative model* in Table 2. Optional breaks of the deterministic term in the periods `t_D1` are treated in accordance with JMN (2000) or KN (2019). The panel extensions of the Johansen procedure are implemented by

```
pcoint.JO(L.data, lags, type, t_D1=NULL, t_D2=NULL, n.factors=FALSE)
pcoint.BR(L.data, lags, type, t_D1=NULL, t_D2=NULL, n.iterations=FALSE)
```

The input argument `L.data` requires a data panel in *list format* as explained above for the argument structure of **pvars** functions in general. `lags` defines the lag order p_i of the individual VAR models in levels and is either a vector of N integers or a single integer for a common lag order $p = p_i$. For assigning the heterogeneous lag orders to each individual, the integers p_i must have the same succession $i = 1, \ldots, N$ as `L.data`. The optional argument `n.factors` can activate the PANIC-defactoring, where the common components with the chosen number of factors $\boldsymbol{F}_t$ are subtracted. Then, the idiosyncratic cointegration rank tests are fixed to the distribution moments for *SL_trend* irrespective of the specification of `type`. Specifically `pcoint.BR()` uses the LM-test from (17) instead, where `n.iterations` defines the number of repetitions in the two-step estimation of β_K. Note that any deterministic term is equipped with a heterogeneous effect β_{0i} in order to comply with the Brownian bridges in the individual $Z_{r_\perp}$. The corresponding option `idx_pool` in `pvarx.VEC()` is thus disabled in `pcoint.BR()` and just cancels out.

The branch of Saikkonen and Lütkepohl. The individual SL-procedure is available via

```
coint.SL(y, dim_p, type_SL, t_D=NULL)
```

The arguments therein are the same as in the individual Johansen procedure except for `type_SL`, which requires a label for the *additive model* from Table 2. Optional breaks of the deterministic trend in the periods `t_D` are treated in accordance with TSL (2008). The panel extensions of the SL procedure are implemented by

```
pcoint.SL(L.data,   lags, type="SL_trend", t_D=NULL, n.factors=FALSE)
pcoint.CAIN(L.data, lags, type="SL_trend", t_D=NULL)
```

Again, the specifications of the input arguments are identical to those of the Johansen procedure `pcoint.JO` except for the `type` of the additive model. The default is `"SL_trend"` as Örsal and Droge (2014) propose for their panel SL-test, Arsova and Örsal (2021) for the CAIN-test, and Örsal and Arsova (2017; 2018) enforce after the defactoring in PANIC.

3.2. Estimating VAR models

The position of the individual VAR model of (1) as the basic econometric unit of Section 2 is reflected in the R implementation by its class `varx`. Accordingly, *(i)* panel VAR estimates of class `pvarx` contain a repeated list `L.varx` of individual VAR, *(ii)* SVAR estimates `id` inherit from the parent class `varx`, and *(iii)* panel SVAR estimates `pid` define their `L.varx` as a list of individual `id` objects. Hence, each individual VAR embedded in the panel estimates can be separately inspected by the methods for `varx` objects.

In **pvars**, `varx` also serves as an intermediary class to ensure compatibility with the other packages of the **vars** ecosystem. Either estimated via **pvars**' `VECM()` or coerced from **vars**' `varest` and `vec2var` objects via `as.varx()`, the `varx` objects can then enter the same functions since the class obeys a unifying construction plan for different VAR model types. In order to apply **pvars** functions to VAR objects of other classes, the user may simply write corresponding `as.varx()` methods instead of altering the original **pvars** function. A list of class `varx` contains the coefficient matrix `$A` for the *full system* and *level representation* of the VAR model, its residual covariance matrix `$SIGMA`, and a structural impact matrix `$B`. In reduced-form VAR objects, the latter is just a placeholder $\texttt{\$B} = I_K$ such that `irf()` generates *forecast-error impulse responses* (Lütkepohl 2005, p. 52). In SVAR objects of class `id`, `$B` is the result of an identification procedure. If a cointegration rank restriction or conditional estimation is employed, the estimates and specifications of these VAR representations are stored in the elements `$VECM`, `$PARTIAL`, and `$MARGINAL` and then transformed to the top-level `$A`.

Panel of VAR models. Extended from the branch of individual VAR resp. VECM, the estimators for the VAR models of heterogeneous panels

```
pvarx.VAR(L.data, lags, type, t_D=NULL, D=NULL)
pvarx.VEC(L.data, lags, dim_r, type, t_D1=NULL, t_D2=NULL,
          D1=NULL, D2=NULL, idx_pool=FALSE, n.iterations=FALSE)
```

use data panels in list format `L.data` to estimate a list of individual VAR models `$L.varx`. The specifications of the VAR processes are the lag orders `lags`, the `type` of the conventional deterministic term, and optional deterministic regressors activated in the periods `t_D` resp. `t_D1` and `t_D2`. While these arguments of `pvarx.VEC()` comply with the labels in Table 2 and with $\boldsymbol{d}_{1,t}$ and $\boldsymbol{d}_{2,t}$ in (2), `pvarx.VAR()` is in accord with **vars**' well-known `VAR()` function and accepts a `"const"`, a linear `"trend"`, `"both"`, or `"none"` in $\boldsymbol{d}_t$ of each individual model in (1). Customized regressors can be included as a common single data matrix or a `list` of individual data matrices (including the presample) via `D` in VAR models and via restricted `D1` and unrestricted `D2` in VECM. Unlike `t_D`, `t_D1`, and `t_D2`, these arguments do not add accompanying lagged regressors to $\boldsymbol{d}_{2,t}$ automatically.

In the next step, the individual coefficients are combined by cross-sectional averages. This provides *(i)* Pesaran and Smith's (1995) mean-group (MG) estimation as suggested by Canova and Ciccarelli (2013) and assessed by Rebucci (2010) for VAR or *(ii)* Pesaran et al.'s (1999) pooled mean-group (PMG) estimation if Breitung's (2005) two-step estimator has been selected. The latter is activated if some variables `idx_pool` are restricted to have homogeneous coefficients in the `dim_r` cointegrating vectors `$beta`. For this, the switching algorithm can be used with further `n.iterations`. If all elements of `idx_pool` are in $[0, \ldots, r]$, the coefficients up to the uniform upper block I_r are throughout heterogeneous and estimated with the individual estimator by Ahn and Reinsel (1990). All resulting panel estimates such as `$A`

or `$beta` are stored in top-level elements of the `pvarx` objects.

3.3. Identifying structure

Equivalently to **svars**' `id.*()` procedures for individual VAR objects, **pvars**' `pid.*()` functions are applied to `pvarx` objects containing the panel estimates of the reduced-form VAR. Accordingly, the arguments to control the underlying identification procedures are identical to those of the **svars** package. Lange *et al.* (2021) describe the implementation and application of the identification procedures in detail.

Imposed. Theoretical considerations like recursive causality may imply restrictions which can be imposed uniformly on B_i of each individual VAR model. Some ensuing functions of Section 3.4 accept a simple list of individual **svars** objects, too, which will produce identical results under these restrictions. However, to enable the full functionality of the **pvars** package, the following functions are added as panel equivalents into the `pid.*()` canon.

```
pid.chol(x, order_k=NULL)
pid.grt(x, LR=NULL, SR=NULL, start=NULL, max.iter=100, conv.crit=1e-07, maxls=1.0)
```

The function `pid.chol()` is the direct extension of **svars**' `id.chol()`, where the optional argument `order_k` allows for specifying alternative causal orderings in the Cholesky decomposition. The panel function `pid.grt()` and its individual counterpart perform the ML estimation of (22) for SVECM under short and long-run restrictions on the $K \times K$ matrices `SR` resp. `LR`. Using the scoring algorithm from **vars**' `SVEC()`, they have identical arguments to tune the optimization procedure as Pfaff (2008b, Sec. 3.2) describes in detail.[19] If the input object `x` contains pooled cointegrating coefficients $\beta_{K,i} = \beta_K$, those are used to calculate the orthogonal complement $\beta_\perp$ in the structural identification of (21).

Data-driven. Residual structure in $\boldsymbol{u}_{it}$ such as non-Gaussianity allows data-driven identification. For this purpose, **pvars** offers the panel applications of ICA. Via the argument `combine`, the user can select a strictly individual identification of B_i as in **svars** (using `"indiv"`), a common rotation of pooled shocks by Herwartz and Wang (2024) (`"pool"`), or `n.factors` common shocks across the individuals by Calhoun *et al.* (2001) (`"group"`).

```
pid.dc(x,  combine, n.factors=NULL, n.iterations=100, PIT=FALSE)
pid.cvm(x, combine, n.factors=NULL, dd=NULL, itermax=500, steptol=100, iter2=75)
```

The panel identification functions pass the combined samples of whitened shocks to the ICA procedures. Like `id.dc()` in **svars**, `pid.dc()` uses the gradient algorithm from **steadyICA** to minimize the distance covariance (dCov) with respect to the rotation angles $\boldsymbol{\theta}$. Their joint option `PIT` activates *probability integral transformation*, which transforms the marginal densities of the structural shocks before evaluating dCov. The maximum number of iterations is `n.iterations=100` by default. The panel function `pid.cvm()` uses the procedure from **svars**' `id.cvm()` to minimize the CvM distance. For both CvM functions, **copula**'s (Kojadinovic and Yan 2010) `indepTestSim()` simulates the distribution of test statistics under independence,

[19] This function integrates **vars**' `SVEC()` into the **pvars** system on the panel level. `SVEC()` cannot be applied to objects of class `cajo.test`, i.e. **urca**'s VECM object with restricted α or β, although these restrictions contribute to the identification of structural shocks. As an individual counterpart, `id.grt()` is applied to `varx` objects, allowing for complex deterministic terms, the MB bootstrap, and `svarirf` methods.

which is either provided via `dd` or called internally if `dd=NULL`. The external provision of `dd` saves computation time if simulated once and then used for multiple calls of `id.cvm()` or `pid.cvm()` on `x` with identical sample dimensions. The remaining arguments control the two-step optimization procedure for $\widehat{\boldsymbol{\theta}}$. In a first step, the differential evolution algorithm from **DEoptim** (Mullen, Ardia, Gil, Windover, and Cline 2011) determines preliminary angles $\boldsymbol{\theta}^*$ within `itermax` iterations under a tolerance of `steptol`. The second step further optimizes the test statistic in `iter2` iterations locally around $\boldsymbol{\theta}^*$.

All `pid.*()` functions extend their `pvarx` input to the class `pid`. Therein, the structural impact matrices $\widehat{\mathsf{B}}_\mathsf{i}$ have been assigned to each SVAR object in `$L.varx` and their mean-group estimates to the top-level element `$B` next to `$A`. By default, the ICA-based panel functions order the columns of all $\widehat{\mathsf{B}}_\mathsf{i}$ uniformly pursuant to the aforementioned convention of the SVAR literature. Consequently, the main diagonal of the mean-group `$B` holds the maximum absolute estimates as positive coefficients $\widehat{\mathsf{b}}_{kk}$. Mean-group statistics of $\widehat{\mathsf{B}}_i$ as presented in Bernoth and Herwartz (2021) and Herwartz and Wang (2024) can be viewed via `pid`'s `summary()` method and exported via `toLatex()`.

3.4. Supporting tools

Dynamic analysis. In the **vars** ecosystem, several tools of dynamic analysis are already available such as *impulse response functions* (IRF), *forecast error variance decomposition* (FEVD) and *historical decomposition* (HD). **pvars** extends this list by *mean-group IRF* (Gambacorta *et al.* 2014, p. 627). Given a VAR object `x` of class `pvarx` or `pid`, the method

```
irf(x, n.ahead=20, normf=NULL, w=NULL)
```

calculates the cross-sectional average of individual responses for each period after the initial impulse. The function optionally provided in `normf` normalizes the shock size of these impulses. Vector `w` with names, N logical, or N numeric elements allows selecting a subset of the N individuals resp. applying real-valued weights in the mean-group estimation.

Persistence profiles (PP) by Pesaran and Shin (1996) are particularly useful for panel cointegration analysis. Given an individual VECM, they map the speed of convergence to the long-run equilibrium after an impulse shock and thus allow counter-checking the individual error correction behavior under a common cointegration rank or pooled cointegration matrix. While a reversion to the r long-run equilibria is the defining property of cointegration, explosive roots can emerge from ignored breaks in the deterministic term on the individual level and contaminate the panel results. In **pvars**, the functions

```
PP.system(x,   n.ahead=20)
PP.variable(x, n.ahead=20, shock=NULL)
```

calculate PP initiated by *system-wide shocks* resp. by *variable-specific shocks* based on the Cholesky decomposition of $\widehat{\Sigma}_{u,i}$. *Structural shocks* are derived from $\widehat{\mathsf{B}}_i$ if `x` is a structural VECM object of class `id` instead of the reduced-form `varx`. The matrix `shock` controls via its K rows which shocks are selected and combined. Both tools, `PP.*()` and `irf()`, return `svarirf` objects, for which **svars** provides the `plot` method to visualize the responses over a horizon of up to `n.ahead` time periods.

Bootstrapping. Bootstrap procedures are a standard tool for VAR modeling to reconstruct the sampling distribution and perform inference. For estimating standard errors of point

estimates and confidence bands of structural impulse responses, **pvars** provides recursive-design bootstrap procedures. In particular, the following functions implement the moving-block bootstrap for individual VAR models (Brüggemann, Jentsch, and Trenkler 2016) and the panel-block bootstrap (Empting *et al.* 2026) respectively.

```
sboot.mb(x, b.length=1, n.ahead=20, n.boot=500, n.cores=1,
         deltas=cumprod((100:0)/100), normf=NULL)
sboot.pmb(x, b.dim=c(1, 1), n.ahead=20, n.boot=500, n.cores=1,
         deltas=cumprod((100:0)/100), normf=NULL, w=NULL)
```

Given an estimated (panel) SVAR object `x` of class `id` resp. `pid`, the bootstrap functions iterate `n.boot` times re-estimating the VAR models, their structural matrices B_i, and impulse responses over a horizon of up to `n.ahead` periods. The arguments from the SVAR object itself (i.e. model specifications, estimation and identification methods, optional restrictions on $\alpha_i\beta^\top$ like rank and weak exogeneity) are passed to and fixed over the bootstrap iterations. In order to speed up their computation by parallel processing, more than one CPU core can be assigned to the procedure via `n.cores`. The resulting **svars** object of class `sboot` allows plotting IRF confidence bands via **svars**' `plot()` method. Confidence intervals for parameters A or B can be viewed via `summary()` and exported via `toLatex()`.

If input `x` contains bias-correction terms `PSI_bc` resp. `L.PSI_bc`, both functions perform a bias-corrected bootstrap. For example, objects from a first bootstrap contain such terms and thus enable the bootstrap-after-bootstrap of individual (Kilian 1998) or panel VAR models (Empting *et al.* 2026), where the weights `deltas` control a successive stationarity correction. `plot()` then displays small-sample-corrected IRF and their confidence bands.

The argument `b.dim` defines the dimensions $\left(b_{(t)}, b_{(i)}\right)$ of the panel blocks for temporal and cross-sectional resampling. The default `c(1, 1)` specifies an *iid* resampling in both dimensions, `c(1, FALSE)` a temporal resampling, and `c(FALSE, 1)` a cross-sectional resampling. Choosing integers $b_{(t)},\ b_{(i)} > 1$ assembles blocks of consecutive residuals to capture residual structure like ARCH or cross-sectional correlation. Moreover, `sboot.mb()` complements **svars**' `mb.boot()` (Lange *et al.* 2021, Sec. 3.6 and 4.2) and accepts individual SVAR objects identified by `id.grt()` (Breitung *et al.* 2004) or `id.iv()` (Jentsch and Lunsford 2022). Here, the default `b.length=1` implies the residual *iid* bootstrap as implemented in **vars**' `irf()`, while a single integer $b_{(t)} > 1$ defines the length of temporal blocks for a moving-block bootstrap.

4. Empirical illustrations

Several empirical illustrations accompany the package to demonstrate its application. In the `help()` for functions, the examples provide chunks of R code for directly copy-pasting unit-tested reproductions. In this section, we focus on the workflow of **pvars** and therefore guide the user to first organize the $K \times T \times N$ data array, then perform a panel cointegration analysis, and finally export the results to LaTeX. Specifically, the reproduced example of Örsal and Arsova (2017) in Section 4.1 illustrates how to perform the *Panel Analysis of Nonstationarity in Idiosyncratic and Common components* (PANIC), and the reproduced example of Arsova and Örsal (2021) in Section 4.2 how to specify deterministic terms. The R code for both illustrations is assembled in the file *pvars_reproductions.R* in **pvars**' *examples* folder.

Data format. The exemplary data sets of the **pvars** package are panels in the prevalent *long format*, where all N multivariate time series have been transposed into $T \times K$ matrices and stacked into an $(N \cdot T) \times (2 + K)$ `data.frame` object. The two additional columns `id_i` and `id_t` contain `factor` elements, which serve as identifiers for individual i and time period t. Accordingly, each observation $\boldsymbol{y}_{it}$ is stored in a single row. The `factor` variables preserve the predefined `levels` order $1, \ldots, N$ within the complete long-format data panel or its subsets.[20] In the following, we consider the data set `MERM` and first extract the names of the variables $k = 1, \ldots, K$ and countries $i = 1, \ldots, N$.

```
R> library("pvars")
R> data("MERM")
R> names_k = colnames(MERM)[-(1:2)]
R> names_i = levels(MERM$id_i)
R> head(MERM, n=3)
```

```
    id_i     id_t          s         m          y         p
1 Brazil 1995_Jan -0.1660546 -3.094546 0.07401953 0.3357538
2 Brazil 1995_Feb -0.1731636 -3.054644 0.07127137 0.3422039
3 Brazil 1995_Mar -0.1176580 -3.055017 0.06986985 0.3539417
```

Naturally, **pvars**' modular implementation works well with panel data in *list format*, where each of the N listed elements is an individual matrix of $T \times K$ time series. This can be constructed by either writing separate time series into the `list` object or transforming the long format data panel via `sapply()`.

```
R> L.data = sapply(names_i, FUN=function(i)
+     ts(MERM[MERM$id_i==i, names_k], start=c(1995, 1), frequency=12),
+     simplify=FALSE)
```

Here, the individual matrices in `L.data` have been defined as time series objects `ts` with `frequency=12` for monthly observations starting in January 1995. Although the functions in **pvars** do not require this, the `ts`-definition simplifies the workflow when using further packages like **ggplot2** (Wickham 2016). The panel functions yet resort to the names of the listed time series as labels for individual results. `sapply()` assigns this definition directly, but `names()` can enforce this subsequently, too, as an alternative transformation requires:

```
R> L.data = lapply(names_i, FUN=function(i) MERM[MERM$id_i==i, names_k])
R> names(L.data) = names_i
```

Either way, the data set is now readily prepared for the econometric analysis with **pvars**.

4.1. The monetary exchange rate model: Conduct a PANIC

[20] Note that `factor()` sorts its `levels` alphabetically unless they are specified explicitly, which applies also to the automatic coercion of `character` columns in R versions prior to 4.0.0. This could lead to mismatches when switching between label standards as in the case of Switzerland with the ISO-3166 abbreviation "CHE".

Örsal and Arsova (2017) illustrate the PANIC analysis of the *monetary exchange rate model* (MERM), where the nominal exchange rate s_{it} between two countries forms a long-run relationship with their relative level of money supply and their relative level of output. As Dąbrowski, Papież, and Śmiech (2014) propose, Örsal and Arsova adopt the log-linear model

$$s_{it} = \mu_{0i} + \mu_{1i}t + \beta_{1i}\left(m_{it} - m_t^*\right) + \beta_{2i}\left(y_{it} - y_t^*\right) + \beta_{3i}\left[\left(p_{it} - p_{it}^T\right) - \left(p_t^* - p_t^{T*}\right)\right] + u_{it}, \quad (36)$$

where the variables for the USA as the preselected reference country are marked with an asterisk. The natural logarithm of the dollar exchange rate for a country i is denoted by s_{it}, the logarithmized nominal money supply by m_{it}, and the logarithmized industrial production index by y_{it}. Moreover, they have included the natural logarithm of consumer price index p_{it} and producer price index p_{it}^T for country i and likewise for the USA.

Data. As `head(MERM)` has shown for the illustrative transformation of the data format, the data set `MERM` contains $K = 4$ variables. These already summarize each log-ratio of (36) and thus enter the additive model in (32) directly as the observed time series $\boldsymbol{y}_{it}$. The monthly observations cover the period 1995/01 – 2007/12 ($T = 156$) for $N = 19$ countries and are listed in `L.data` after transforming their data format. The names of `L.data`'s 19 elements provide the labels for the "individuals" in Table 4.

Approximate factor model. The first step of PANIC is to estimate the approximate factor model in (32), which splits $\boldsymbol{y}_{it}$ into common and idiosyncratic components $\Lambda_i^\top \boldsymbol{F}_t$ resp. $\boldsymbol{y}_{it}^{id}$. The factor model considers the data panel just as a collection of time series without individual structure. Both dimensions $T \times NK$ of the data are assumed to be large and both components of the model may involve mixes of $I(0)$ and $I(1)$ series. First-differencing these data panels beforehand is a valid choice to estimate the factor model by PCA (Bai and Ng 2004) and to determine its number of common factors $\boldsymbol{F}_t$ by the eigenvalues.[21] The information criteria in `[[1]]` however ignore the individual structure of our panel and thus tend to pick up the domestic dependencies between the $K = 4$ variables within countries. Since we are interested in the factors describing cross-sectional dependence only, we prefer the specification procedure by Onatski (2010). His *edge distribution* `ED` is more robust against domestic dependencies because it looks for a characteristic kink in the ordered eigenvalues. `ED` also works with the original `L.data` in levels irrespective of the components' order of integration. In order to find the optimal number of factors within the discrete interval $\{0, \ldots, 20\}$, we enter the R function

```
R> speci.factors(L.data, k_max=20, n.iterations=4)

### Optimal number of common factors ###
[[1]]
   PC IC IPC
p1 20 20   7
p2 20 20   7
p3 20 20   5

[[2]]
ER GR ED
1  2  8
```

[21] See Corona, Poncela, and Ruiz (2017) for an overview and Monte Carlo results. An exception is the set of $IPC(k)$ criteria by Bai (2004), who seeks to distinguish nonstationary factors from stationary idiosyncratic series. Accordingly, `speci.factors()` suppresses their result if `differenced=TRUE` is selected.


A numerical result for `ED` indicates that the default of `n.iterations=4` allows Onatski's (2010) edge distribution to converge. In case of an `NA`, the user needs to increase the number of iterations, but small numbers are often sufficient.

The result of eight common factors can be visualized and checked in a scree plot. In accordance with PANIC of the `pcoint` functions, we may consider the factor model estimated with the first-differenced and standardized data now:

```
R> R.fac0 = speci.factors(L.data, k_max=20, n.iterations=4,
+    differenced=TRUE, centered=TRUE, scaled=TRUE, n.factors=8)

R> library("ggplot2")
R> pal = c("#999999", RColorBrewer::brewer.pal(n=8, name="Spectral"))
R> lvl = levels(R.fac0$eigenvals$scree)
R> ggplot(R.fac0$eigenvals[1:20, ]) +
+    geom_col(aes(x=n, y=share, fill=scree), color="black", width=0.75) +
+    scale_fill_manual(values=pal, breaks=lvl, guide="none") +
+    labs(x="Component number", y="Share of total variance", title=NULL) +
+    theme_bw()
```

Figure 3: Scree plot.



In the resulting plot of Figure 3, the first eight eigenvalues for the relevant components are colored.[22] They still account for almost 50% of the total variation in the first-differenced and centered $K \cdot N$ time series, but the PCA of the original data attributes over 98% to the first principal component alone. It appears that this all-dominating component is a linear trend in the time series, which is removed after first-differencing and centering. Now, the eighth and ninth eigenvalue are inconsiderable, while the first two exhibit more pronounced kinks. Indeed, the *eigenvalue ratios* `ER` and *growth rates* `GR` by Ahn and Horenstein (2013) as well as `ED` by Onatski (2010) hint at one resp. two common factors.

```
R> R.fac0$selection[[2]]

ER GR ED
1  1  2
```

[22] The vector graphics in this LaTeX document have been generated by the **tikzDevice** package (Sharpsteen and Bracken 2020), which prints R plots as a Ti*k*Z environment into ".tex" files.

For the reproduction, we yet proceed with the decision by Örsal and Arsova (2017) and adhere to the conservative choice of eight common factors in order to ensure cross-sectional independence for the panel test.

Panel cointegration tests. The approximate factor model with `n.factors=8` yields a nonstationary,[23] idiosyncratic remainder $\hat{\boldsymbol{y}}_{it}^{id}$, to which Örsal and Arsova (2017) apply the panel SL-tests. To reproduce their results, we specify **pvars**' function `pcoint.SL()` as follows. Due to the defactoring and in accordance with the econometric model in (36), the $N = 19$ individual testing procedures therein must take care of deterministic trends. The lag order p_i of each idiosyncratic VAR model is chosen from the discrete interval $\{1, \ldots, 4\}$ by the minimized *Akaike information criterion*. Here, we enter the results directly, but **vars** functions may determine them from the data matrices in `R.fac0$L.idio`, too.

```
R> R.lags = c(2, 2, 2, 2, 1, 2, 2, 4, 2, 3, 2, 2, 2, 2, 2, 1, 1, 2, 2)
R> R.pcsl = pcoint.SL(L.data, lags=R.lags, type="SL_trend", n.factors=8)
R> toLatex(R.pcsl)
```

The method `toLatex()` prints the `pcoint` results as a `tabular` for LaTeX, which has been encapsulated in LaTeX's float environment in order to create Table 4. The table reports the

Table 4: Panel cointegration rank tests for MERM.

Individual		statistics				p-values			
	lags	$r_{H0} = 0$	$r_{H0} = 1$	$r_{H0} = 2$	$r_{H0} = 3$	$r_{H0} = 0$	$r_{H0} = 1$	$r_{H0} = 2$	$r_{H0} = 3$
Brazil	2	41.606	13.963	8.660	3.618	0.115	0.832	0.473	0.256
Canada	2	43.889	19.816	6.253	0.413	0.070	0.406	0.747	0.940
Colombia	2	25.210	13.042	4.744	1.914	0.875	0.881	0.891	0.558
Czech Republic	2	29.238	17.663	7.712	0.583	0.689	0.568	0.580	0.901
Denmark	1	37.925	18.804	7.758	2.450	0.230	0.480	0.575	0.442
Hungary	2	32.372	16.940	8.401	0.863	0.510	0.624	0.502	0.829
India	2	24.846	14.801	6.280	1.889	0.887	0.780	0.744	0.564
Indonesia	4	26.911	12.640	4.211	1.687	0.806	0.899	0.928	0.612
Israel	2	36.282	21.678	6.217	0.561	0.301	0.285	0.751	0.906
Japan	3	28.154	15.395	6.966	1.975	0.746	0.739	0.667	0.544
Korea	2	57.469	13.816	7.835	2.817	0.002	0.840	0.566	0.374
Mexico	2	28.996	20.596	6.638	0.682	0.702	0.352	0.704	0.876
Norway	2	43.766	19.633	6.715	1.367	0.072	0.419	0.696	0.694
Poland	2	60.457	28.641	6.618	0.557	0.001	0.048	0.707	0.907
South Africa	2	21.298	10.053	6.868	4.730	0.968	0.976	0.678	0.147
Sweden	1	32.127	7.147	3.386	0.777	0.524	0.998	0.969	0.851
Switzerland	1	28.419	13.682	3.413	1.699	0.733	0.848	0.968	0.609
Turkey	2	48.692	27.137	14.041	0.325	0.021	0.075	0.094	0.958
United Kingdom	2	50.253	24.440	4.287	3.031	0.014	0.152	0.923	0.339
Panel		statistics				p-values			
		$r_{H0} = 0$	$r_{H0} = 1$	$r_{H0} = 2$	$r_{H0} = 3$	$r_{H0} = 0$	$r_{H0} = 1$	$r_{H0} = 2$	$r_{H0} = 3$
LRbar		2.305	-1.346	-2.635	-2.095	0.011	0.911	0.996	0.982
Choi P		70.515	29.377	17.050	20.338	0.001	0.841	0.999	0.992
Choi Pm		3.730	-0.989	-2.403	-2.026	0.000	0.839	0.992	0.979
Choi Z		-1.914	1.528	2.639	2.124	0.028	0.937	0.996	0.983

individual and panel results for each hypothesis $r_{H0} = 0, \ldots, K-1$, which refer to Table 5 in Örsal and Arsova (2017, p. 68). All four combination approaches under the independence assumption of the idiosyncratic VAR processes have been used. Comparing the p-values to a significance level of $\alpha = 5\%$, all sequential panel test procedures reject the hypotheses up to $r_{H0} = 1$ and thus confirm the presence of a single cointegration relation in $\boldsymbol{y}_{it}^{id}$.

[23] Single-equation PANIC methods, i.e. unit root and residual-based cointegration tests, are implemented in **PANICr** by Bronder (2016). The R package was archived on CRAN in 2020.

Cointegration rank of the factors. Having determined the idiosyncratic cointegration rank, the PANIC turns then to the cointegration within the eight common factors $\boldsymbol{F}_t$. The `$CSD` element of the `pcoint` object contains the estimates for the cross-sectional dependence and is identical for the PANIC analysis of any `pcoint` function. Therein, the eigenvalues of the PCA are stored in the vector `eigenvals` and the cumulated common factors in the matrix `Ft` of dimension `dim_T` $\times$ `n.factors`. These multivariate time series shall be plotted first to get an overview as in Figure 4 of Örsal and Arsova (2017, p. 71). For this, we define the factor matrix `Ft` as a `ts` object with the same specifications as the observed time series and use the related plotting method via `autoplot()`. The package **ggfortify** (Tang, Horikoshi, and Wenxuan 2016) provides a comprehensive set of unified methods for **ggplot2** graphics.

```
R> library("ggfortify")
R> Ft = ts(R.pcsl$CSD$Ft, start=c(1995, 1), frequency=12)
R> autoplot(Ft, facets=FALSE, size=1.75) + theme_bw() +
+     scale_color_brewer(palette="Spectral") +
+     labs(x=NULL, y=NULL, color="Factor", title=NULL)
```

Figure 4: Estimated common factors.




Figure 4 depicts the `autoplot()` result for the common factors, which are ordered according to their decreasing eigenvalue of the PCA. Using `palette="Spectral"` for factors' color assignment, the transition from "warm" to "cold" colors reflects decreasing power of the factors to explain the common variation within the observed time series. In comparison to the figure from the original study, the factors $\hat{\boldsymbol{F}}_t$ may have switched signs and be multiplied by a scalar such that only their dynamics are informative. However, the PANIC analysis of both components is invariant to this differing normalization of loadings and factors $\hat{\Lambda}_i^\top \hat{\boldsymbol{F}}_t$: *(1)* The idiosyncratic components are calculated via this complete term of common components, which the normalization does not alter. *(2)* The common factors $\hat{\boldsymbol{F}}_t$ exhibit the same strength of cointegration, i.e. $\hat{\boldsymbol{\lambda}}$ of the reduced-rank regression in (8), irrespective of any scaling weight multiplied to a time series of $\hat{\boldsymbol{F}}_t$.

In order to determine the cointegration rank within $\boldsymbol{F}_t$, we employ the single VECM in (2) with a linear trend in the cointegration space. The proceeding is the same as any individual analysis of the country-specific time series. The AIC, as provided by **vars**' `VARselect()` or **pvars**' `speci.VAR()`, hints at a lag order of $p_{F_t} = 2$ for the VAR model in (1) without rank-restriction. We perform both procedures, Johansen (1996) and SL (2000c), using an

unrestricted intercept and a linear trend restricted to the cointegration relations (*Case 4*, see Table 2) and the corresponding intercept and linear trend in the additive (10) (*SL_trend*).

```
R> R.Ftsl = coint.SL(R.pcsl$CSD$Ft, dim_p=2, type_SL="SL_trend")
R> R.Ftjo = coint.JO(R.pcsl$CSD$Ft, dim_p=2, type="Case4")
R> toLatex(R.Ftsl, R.Ftjo, write_ME=TRUE,
+     add2header=c("\\textbf{SL:} Trace test",
+                  "\\textbf{Johansen:} Trace test"))
```

Both results are jointly exported to LaTeX via the `toLatex()` method for `coint` objects. In order to distinguish them in the joint table, the optional argument `add2header` labels the procedure in the respective column. Depending on the input `coint` objects, the user could indicate individual names or different specifications manually here as well. In contrast to the panel test functions, the two functions of the `coint` branches conduct the maximum eigenvalue LR-test additionally. The argument `write_ME` controls whether these results shall enter the LaTeX table. This feature is particularly useful for test procedures on more complex data-generating processes with trend breaks or weakly exogenous variables. In these cases, the distribution moments of $Z_{r_\perp}$ and thus p-values would be available for the trace test only such that the element `$pvals_ME` of the `coint` object would report a vector of `NA`s. However, since the empirical example implies standard specifications, the complete Table 5 has been printed with `write_ME=TRUE`.

Table 5: Cointegration rank tests for the common factors.

	SL: Trace test		Max. eigenvalue test		**Johansen:** Trace test		Max. eigenvalue test	
r_{H0}	statistic	p-value	statistic	p-value	statistic	p-value	statistic	p-value
0	202.448	0.000	63.178	0.001	251.612	0.000	74.245	0.000
1	116.540	0.080	40.822	0.124	177.367	0.000	57.414	0.006
2	85.131	0.125	38.750	0.043	119.952	0.034	38.361	0.206
3	47.891	0.622	20.551	0.637	81.591	0.147	30.719	0.297
4	27.169	0.794	14.439	0.716	50.873	0.379	19.469	0.699
5	14.737	0.784	10.300	0.642	31.404	0.427	14.536	0.684
6	3.719	0.955	3.465	0.894	16.868	0.433	9.713	0.656
7	1.406	0.684	1.406	0.684	7.155	0.338	7.155	0.339

The table shows results for the four tests on the cointegration rank within the eight common factors. In accordance with Table 6 in Örsal and Arsova (2017, p. 68), the trace tests in the SL- and the Johansen procedure stop at an r_{H0} of one resp. three cointegration relations and thus suggest at least five global stochastic trends driving the observable variables $\boldsymbol{y}_{it}$ in the PANIC model of (32). Correspondingly, the maximum eigenvalue test procedures suggest ranks of one resp. two at a significance level of 5%. The range of five to seven global stochastic trends is also in accord with the $IPC(k)$ criteria.

4.2. The exchange rate pass-through: Specify the deterministic term

The empirical example from Arsova and Örsal (2016, Ch. 6) about the *exchange rate pass-through* (ERPT) shows how to perform cointegration rank analysis under the consideration of structural breaks in the cointegration relation. Their example itself is based on the data set and theoretical model from Banerjee and Carrion-i-Silvestre (2015), who assess a single long-run relationship between the logarithmized time series of import price mp_{it}, exchange

rate er_{it}, and foreign price fp_t in the linear model

$$mp_{it} = \mu_{0i} + \mu_{1i}t + \beta_{1i} \cdot er_{it} + \beta_{2i} \cdot fp_t + u_{it}. \tag{37}$$

Data.[24] The data set on the $K = 3$ variables $\boldsymbol{y}_{it}$ consists of monthly observations over the period 1995/01 – 2005/03 ($T = 123$) for $N = 7$ Euro-area countries and nine different industries. However for the illustration, we reproduce the empirical analysis for the industry *"chemicals and related products"* only and it is up to the user to try out cointegration tests for the remaining industries. Their time series are also stored in **pvars**' data set `ERPT`. In order to subset and transform the long-format panel `ERPT` into the necessary list format, we follow the same steps as explained at the beginning of Section 4 and select the variables for the chemical industry denoted by serial number 5. Only for an easier comparison with the results from Arsova and Örsal (2016), the country names `names_i` are reordered according to the original output tables. This has no consequences for the implementation.

```
R> library("pvars")
R> data("ERPT")
R> names_k = c("lpm5", "lfp5", "llcusd")
R> names_i = levels(ERPT$id_i)[c(1,6,2,5,4,3,7)]
R> L.data  = sapply(names_i, FUN=function(i)
+     ts(ERPT[ERPT$id_i==i, names_k], start=c(1995, 1), frequency=12),
+     simplify=FALSE)
```

Over the considered sample period, Arsova and Örsal (2016) suspect a level shift and trend break in May 2002 motivated by the appreciation of the Euro against the US dollar. The authors attribute this persistent change to effects from the outside of the ERPT model in (37), namely *(i)* the aftermath of the terrorist attacks on the World Trade Center on 11 September 2001, *(ii)* hesitant investors on the US markets, *(iii)* the declining importance of US exports on the world markets, and *(iv)* the fully established Euro after the national currencies had been withdrawn from circulation in March 2002. Since the complete data set including the presample periods comprises the time interval 1995/01 – 2005/03, the single break period τ of 2002/05 is counted to be the 89th one within the interval $1, \ldots, 123$.

Specify the deterministic term. In order to accommodate this structural break, the CAIN-test considers individual TSL-procedures by Trenkler *et al.* (2008), which allow for individual-specific deterministic components in the additive (10) given by

$$M_{\mu,i}\boldsymbol{d}_{it} = [\boldsymbol{\mu}_{0i} : \boldsymbol{\mu}_{1i} : \boldsymbol{\delta}_{0i} : \boldsymbol{\delta}_{1i}]\, \boldsymbol{d}_{it} = \boldsymbol{\mu}_{0i} + \boldsymbol{\mu}_{1i}t + \boldsymbol{\delta}_{0i}d_{it} + \boldsymbol{\delta}_{1i}b_{it}. \tag{38}$$

Besides the conventional constant 1 and linear trend t, the $n_{d,i} \times 1$ vectors $\boldsymbol{d}_{it}$ stack the period-specific shift dummies d_{it} and trend breaks b_{it}. These regressors are $d_{it} = b_{it} = 0$ at first and activated to be $d_{it} = 1$ and $b_{it} = t - \tau_i + 1$ when $t \geq \tau_i$. The empirical example holds the special characteristics that there is only a single break and it occurs at the very same period $\tau_i = \tau = 89$ in each i. Therefore, the shift, which must accompany the single break, is the sole $d_t = \Delta b_t$ and the deterministic regressors $\boldsymbol{d}_t = (1, t, \Delta b_t, b_t)^\top$ are in fact identical for each i. This has no further consequences for the flexible CAIN-TSL test itself,

[24] The balanced panel `ERPT` as used by Arsova and Örsal (2016) contains fewer individuals than Banerjee and Carrion-i-Silvestre (2015) actually provide. The countries Austria, Finland, and Portugal are omitted because Eurostat as the primary data source has reported some missing values in these time series.

but its implementation `pcoint.CAIN()` then requires only the single value $\tau = 89$ instead of a list of N specifications for the argument `t_D`. Accordingly, the objects `R.t_D` and `L.t_D` lead to identical results. The latter can serve as a basis for adding individual-specific regressors d_{it}. In this example, the explicit formulation for France is yet redundant as **pvars** adds any accompanying shift from `t_break` automatically.

```
R> R.t_D = list(t_break=89)
R> L.t_D = sapply(names_i, function(i) list(t_break=89), simplify=FALSE)
R> L.t_D$France$t_shift = c(89)
```

For the feasible GLS-detrending,[25] Table 2 shows how **pvars** constructs $\boldsymbol{d}_{1,it}$ and $\boldsymbol{d}_{2,it}$ in the preceding reduced-rank regressions of $\Delta \boldsymbol{y}_{it}$ on $\boldsymbol{z}_{1,it} = \left(\boldsymbol{y}_{i,t-1}^\top, \boldsymbol{d}_{1,it}^\top\right)^\top$ corrected for $\boldsymbol{z}_{2,it} = \left(\Delta \boldsymbol{y}_{i,t-1}^\top, \ldots, \Delta \boldsymbol{y}_{i,t-p_i+1}^\top, \boldsymbol{d}_{2,it}^\top\right)^\top$. The deterministic regressors $\boldsymbol{d}_{1,it} = (t-1, b_{t-1})^\top$ for the cointegration relations consist of a linear trend and its break in $\tau = 89$. In addition to the unrestricted constant, **pvars** includes the shift Δb_t and lags of the impulse dummy $d_{\tau,t}^{\text{im}}$ into $\boldsymbol{d}_{2,it} = \left(1, \Delta b_t, d_{\tau,t}^{\text{im}}, \ldots, d_{\tau,t-(p_i-1)}^{\text{im}}\right)^\top$ automatically. In converse notation, these lags can be easily recognized as a solitary impulse $d_{\bullet,t}^{\text{im}} = 1$ in each period $\bullet \in \{\tau, \ldots, \tau + (p_i - 1)\}$.

Panel cointegration tests. Arsova and Örsal (2016) determine the individual lag orders p_i with the modified AIC by Qu and Perron (2007) under the null hypothesis of no cointegration. The feasible estimation of $M_{\mu,i}$ in (38) and the LR-test based on an individual VECM with detrended time series then proceed as generally described in Section 2.1. Finally, the panel statistics of (33) by Hartung (1999) and Arsova and Örsal (2021) combine the individual p-values under the three correction factors to account for cross-sectional dependence. We conduct the complete CAIN test procedure with the `pcoint` command

```
R> R.lags = c(3, 3, 3, 4, 4, 3, 4)  # by modified AIC
R> R.cain = pcoint.CAIN(L.data, lags=R.lags, t_D=R.t_D, type="SL_trend")
R> R.cain$CSD$rho_tilde
```

```
            r_H0 = 0  r_H0 = 1  r_H0 = 2
Hartung_K1 0.8123894 0.3973220 0.5726069
Hartung_K2 0.7954536 0.3583592 0.5403518
CAIN       0.1252727 0.1310724 0.1448498
```

```
R> R.cain$CSD$rho_eps
```

```
[1] 0.6333148
```

In the element `$CSD`, the matrix `rho_tilde` contains the correction factors for each panel test and hypothesis $r_{H0} = 0, \ldots, K-1$. Further, `rho_eps` stores the *average absolute cross-sectional correlation* ρ_ϵ calculated only for $r_{H0} = 0$. Its high value indicates strong cross-sectional dependence for the empirical example. Arsova and Örsal (2016, p. 15) attribute this to the common foreign price series fp_t and similar exchange rate dynamics er_{it}. After the Euro exchange rate was fixed on 31 December 1998, the variable er_{it} has been identical across *(i)* six of the $N = 7$ European countries since January 1999 and *(ii)* Greece subsequently since its adoption of the euro in January 2001.

[25] See TSL (2008, p. 335) for more details and Arsova and Örsal (2021, Ch. 6) for another panel example.

```
R> toLatex(R.cain)
```

Table 6: Panel cointegration rank tests for ERPT.

Individual		statistics			p-values		
	lags	$r_{H0} = 0$	$r_{H0} = 1$	$r_{H0} = 2$	$r_{H0} = 0$	$r_{H0} = 1$	$r_{H0} = 2$
France	3	35.006	13.709	3.812	0.024	0.248	0.425
Netherlands	3	32.854	11.754	5.514	0.044	0.402	0.220
Germany	3	36.446	20.365	2.438	0.015	0.029	0.666
Italy	4	39.369	18.309	1.245	0.006	0.060	0.888
Ireland	4	32.834	19.624	1.814	0.045	0.038	0.786
Greece	3	34.559	16.713	3.900	0.027	0.102	0.412
Spain	4	28.230	9.743	2.108	0.146	0.598	0.730
Panel		statistics			p-values		
		$r_{H0} = 0$	$r_{H0} = 1$	$r_{H0} = 2$	$r_{H0} = 0$	$r_{H0} = 1$	$r_{H0} = 2$
Hartung $K1$		-2.034	-1.477	0.335	0.021	0.070	0.631
Hartung $K2$		-2.052	-1.531	0.343	0.020	0.063	0.634
CAIN		-3.725	-2.033	0.517	0.000	0.021	0.697

As in Section 4.1, the `pcoint` object is printed by `toLatex()` and encapsulated in LaTeX's `table` environment. Table 6 displays the individual and panel results for each hypothesis $r_{H0} = 0, \ldots, K-1$, which Arsova and Örsal (2016, p. 22) report in Tables 7 and 8 for the industry of *"5: Chemicals and related products"*, too. Comparing the p-values to a significance level of $\alpha = 5\%$, the sequential testing procedure of CAIN rejects the hypotheses up to $r_{H0} = 2$ and thus suggests the presence of two cointegration relations overall.

5. Summary

This article has presented a set of VAR methods for panel data (Section 2), described their implementation in **pvars** (Section 3), and illustrated their application (Section 4). The R package comprises *panel cointegration rank tests* as well as *estimators of VAR models for heterogeneous panels* and *panel methods for structural identification* of the reduced-form VAR models. In this context, **pvars** addresses typical properties of financial and macroeconomic panel data, in particular cross-sectional dependence and structural breaks in the deterministic term. Finally, **pvars** supplements functions for model specification and dynamic analysis which are not provided by other packages of the **vars** ecosystem, namely various *criteria for the number of common factors*, *persistence profiles*, *mean-group IRF*, and *moving-block bootstrap procedures* for panel SVAR models.

References

Abrigo M, Love I (2016). "Estimation of Panel Vector Autoregression in Stata." *Stata Journal*, **16**(3), 778–804. doi:10.1177/1536867X1601600314.

Ahn SK, Horenstein A (2013). "Eigenvalue Ratio Test for the Number of Factors." *Econometrica*, **81**(3), 1203–1227. doi:10.3982/ECTA8968.

Ahn SK, Reinsel GC (1990). "Estimation for Partially Nonstationary Multivariate Autoregressive Models." *Journal of the American Statistical Association*, **85**(411), 813–823. doi:10.1080/01621459.1990.10474945.

Amisano G, Giannini C (1997). *Topics in Structural VAR Econometrics*. 2nd edition. Springer. ISBN 978-3-642-60623-6.

Anderson TW (1951). "Estimating Linear Restrictions on Regression Coefficients for Multivariate Normal Distributions." *Annals of Mathematical Statistics*, **22**(3), 327–351. doi:10.1214/aoms/1177729580.

Anderson TW (2003). *An Introduction to Multivariate Statistical Analysis*. Wiley Series in Probability and Statistics, 3rd edition. Wiley. ISBN 0471360910,9780471360919.

Arsova A, Örsal DDK (2016). "A Panel Cointegrating Rank Test with Structural Breaks and Cross-Sectional Dependence." *Beiträge zur Jahrestagung des Vereins für Socialpolitik 2016: Demographischer Wandel - Session: Time Series Econometrics No. D01-V3*, German Economic Association.

Arsova A, Örsal DDK (2018). "Likelihood-based Panel Cointegration Test in the Presence of a Linear Time Trend and Cross-Sectional Dependence." *Econometric Reviews*, **37**(10), 1033–1050. doi:10.1080/07474938.2016.1183070.

Arsova A, Örsal DDK (2021). "A Panel Cointegrating Rank Test with Structural Breaks and Cross-Sectional Dependence." *Econometrics and Statistics*, **17**(C), 107–129. doi:10.1016/j.ecosta.2020.05.002.

Bai J (2004). "Estimating Cross-Section Common Stochastic Trends in Nonstationary Panel Data." *Journal of Econometrics*, **122**(1), 137–183. doi:10.1016/j.jeconom.2003.10.022.

Bai J, Ng S (2002). "Determining the Number of Factors in Approximate Factor Models." *Econometrica*, **70**(1), 191–221. doi:10.1111/1468-0262.00273.

Bai J, Ng S (2004). "A PANIC Attack on Unit Roots and Cointegration." *Econometrica*, **72**(4), 1127–1177. doi:10.1111/j.1468-0262.2004.00528.x.

Bai J, Perron P (1998). "Estimating and Testing Linear Models with Multiple Structural Changes." *Econometrica*, **66**(1), 47–78. doi:10.2307/2998540.

Bai J, Perron P (2003). "Computation and Analysis of Multiple Structural Change Models." *Journal of Applied Econometrics*, **18**(1), 1–22. doi:10.1002/jae.659.

Bakirov NK, Rizzo ML, Székely GJ (2006). "A Multivariate Nonparametric Test of Independence." *Journal of Multivariate Analysis*, **97**(8), 1742–1756. doi:10.1016/j.jmva.2005.10.005.

Banerjee A, Carrion-i-Silvestre JL (2015). "Cointegration in Panel Data with Structural Breaks and Cross-Section Dependence." *Journal of Applied Econometrics*, **30**(1), 1–23. doi:10.1002/jae.2348.

Bernoth K, Herwartz H (2021). "Exchange Rates, Foreign Currency Exposure and Sovereign Risk." *Journal of International Money and Finance*, **117**(C), 102454. doi:10.1016/j.jimonfin.2021.102454.

Blackburne E, Frank M (2007). "Estimation of Nonstationary Heterogeneous Panels." *Stata Journal*, **7**(2), 197–208. doi:10.1177/1536867X0700700204.

Boeck M, Feldkircher M, Huber F (2022). "BGVAR: Bayesian Global Vector Autoregressions with Shrinkage Priors in R." *Journal of Statistical Software*, **104**(9), 1–28. doi:10.18637/jss.v104.i09.

Breitung J (2005). "A Parametric Approach to the Estimation of Cointegration Vectors in Panel Data." *Econometric Reviews*, **24**(2), 151–173. doi:10.1081/ETC-200067895.

Breitung J, Brüggemann R, Lütkepohl H (2004). "Structural Vector Autoregressive Modeling and Impulse Responses." In H Lütkepohl, M Krätzig (eds.), *Applied Time Series Econometrics*, Themes in Modern Econometrics, chapter 4, pp. 159–196. Cambridge University Press. ISBN 9780521547871.

Bronder S (2016). "**PANICr**: Panel Analysis of Nonstationarity in Idiosyncratic and Common Components." *R package version 1.0*, CRAN. URL https://CRAN.R-project.org/package=PANICr.

Brüggemann R, Jentsch C, Trenkler C (2016). "Inference in VARs with Conditional Heteroskedasticity of Unknown Form." *Journal of Econometrics*, **191**(1), 69–85. doi:10.1016/j.jeconom.2015.10.004.

Brüggemann R, Lütkepohl H (2005). "Practical Problems with Reduced-Rank ML Estimators for Cointegration Parameters and a Simple Alternative." *Oxford Bulletin of Economics and Statistics*, **67**(5), 673–690. doi:10.1111/j.1468-0084.2005.00136.x.

Calhoun VD, Adali T, Pearlson GD, Pekar JJ (2001). "A Method for Making Group Inferences from Functional MRI Data Using Independent Component Analysis." *Human Brain Mapping*, **14**(3), 140–151. doi:10.1002/hbm.1048.

Canova F, Ciccarelli M (2013). "Panel Vector Autoregressive Models: A Survey." *Advances in Econometrics*, **32**, 205–246. doi:10.1108/S0731-9053(2013)0000031006.

Carrion-i Silvestre JL, Surdeanu L (2011). "Panel Cointegration Rank Testing with Cross-Section Dependence." *Studies in Nonlinear Dynamics & Econometrics*, **15**(4), 1–43. doi:10.2202/1558-3708.1825.

Choi I (2001). "Unit Root Tests for Panel Data." *Journal of International Money and Finance*, **20**(2), 249–272. doi:10.1016/S0261-5606(00)00048-6.

Christopoulos D, Tsionas M (2004). "Financial Development and Economic Growth: Evidence from Panel Unit Root and Cointegration Tests." *Journal of Development Economics*, **73**(1), 55–74. doi:10.1016/j.jdeveco.2003.03.002.

Comon P (1994). "Independent Component Analysis, A New Concept?" *Signal Processing*, **36**, 287–314. doi:10.1016/0165-1684(94)90029-9.

Corona F, Poncela P, Ruiz E (2017). "Determining the Number of Factors after Stationary Univariate Transformations." *Empirical Economics*, **53**(1), 351–372. doi:10.1007/s00181-016-1158-5.

Croissant Y, Millo G (2008). "Panel Data Econometrics in R: The **plm** Package." *Journal of Statistical Software*, **27**(2). doi:10.18637/jss.v027.i02.

Doornik J (1998). "Approximations to the Asymptotic Distributions of Cointegration Tests." *Journal of Economic Surveys*, **12**(5), 573–93. doi:10.1111/1467-6419.00068.

Dąbrowski MA, Papież M, Śmiech S (2014). "Exchange Rates and Monetary Fundamentals in CEE Countries: Evidence from a Panel Approach." *Journal of Macroeconomics*, **41**(C), 148–159. ISSN 0164-0704. doi:10.1016/j.jmacro.2014.05.005.

Empting LFT, Herwartz H (2025). "Revisiting the 'Productivity of Public Capital': VAR Evidence on the Heterogeneous Dynamics in a New Panel of 23 OECD Countries." *Working paper*, Universität Göttingen.

Empting LFT, Öztürk S, Maxand S, Wagner K (2026). "Inference in Panel SVARs with Two-Way Dependence." arXiv:2610.11898 [econ.EM]. doi:10.48550/arXiv.2610.11898.

Fisher RA (1932). *Statistical Methods for Research Workers*, volume 5 of *Biological Monographs and Manuals*. Oliver and Boyd.

Gambacorta L, Hofmann B, Peersman G (2014). "The Effectiveness of Unconventional Monetary Policy at the Zero Lower Bound: A Cross-Country Analysis." *Journal of Money, Credit and Banking*, **46**(4), 615–642. doi:10.1111/jmcb.12119.

Genest C, Quessy JF, Rémillard B (2007). "Asymptotic Local Efficiency of Cramér–von Mises Tests for Multivariate Independence." *Annals of Statistics*, **35**(1), 166–191. doi:10.1214/009053606000000984.

Groen JJ, Kleibergen F (2003). "Likelihood-based Cointegration Analysis in Panels of Vector Error-Correction Models." *Journal of Business & Economic Statistics*, **21**(2), 295–318. doi:10.1198/073500103288618972.

Hamilton JD (1994). *Time Series Analysis*. Princeton University Press. ISBN 0691042896.

Hartung J (1999). "A Note on Combining Dependent Tests of Significance." *Biometrical Journal*, **41**(7), 849–855. doi:10.1002/(SICI)1521-4036(199911)41:7<849::AID-BIMJ849>3.0.CO;2-T.

Herwartz H (2018). "Hodges–Lehmann Detection of Structural Shocks: An Analysis of Macroeconomic Dynamics in the Euro Area." *Oxford Bulletin of Economics and Statistics*, **80**(4), 736–754. doi:10.1111/obes.12234.

Herwartz H, Lange A, Maxand S (2022). "Data-Driven Identification in SVARs – When and How Can Statistical Characteristics Be Used to Unravel Causal Relationships?" *Economic Inquiry*, **60**(2), 668–693. doi:10.1111/ecin.13035.

Herwartz H, Wang S (2024). "Statistical Identification in Panel Structural Vector Autoregressive Models Based on Independence Criteria." *Journal of Applied Econometrics*, **39**(4), 620–639. doi:10.1002/jae.3044.

Hlouskova J, Wagner M (2010). "The Performance of Panel Cointegration Methods: Results from a Large Scale Simulation Study." *Econometric Reviews*, **29**(2), 182–223. doi:10.1080/07474930903382182.

Hodges JL, Lehmann EL (1963). "Estimates of Location Based on Rank Tests." *Annals of Mathematical Statistics*, **34**(2), 598–611. doi:10.1214/aoms/1177704172.

Im KS, Pesaran MH, Shin Y (2003). "Testing for Unit Roots in Heterogeneous Panels." *Journal of Econometrics*, **115**(1), 53–74. doi:10.1016/S0304-4076(03)00092-7.

Izenman AJ (1975). "Reduced-Rank Regression for the Multivariate Linear Model." *Journal of Multivariate Analysis*, **5**(2), 248–264. doi:10.1016/0047-259X(75)90042-1.

Jentsch C, Lunsford KG (2022). "Asymptotically Valid Bootstrap Inference for Proxy SVARs." *Journal of Business & Economic Statistics*, **40**(4), 1876–1891. doi:10.1080/07350015.2021.1990770.

Johansen S (1988). "Statistical Analysis of Cointegration Vectors." *Journal of Economic Dynamics & Control*, **12**(2-3), 231–254. doi:10.1016/0165-1889(88)90041-3.

Johansen S (1991). "Estimation and Hypothesis Testing of Cointegration Vectors in Gaussian Vector Autoregressive Models." *Econometrica*, **59**(6), 1551–1580. ISSN 00129682, 14680262. doi:10.2307/2938278.

Johansen S (1996). *Likelihood-based Inference in Cointegrated Vector Autoregressive Models*. Advanced Texts in Econometrics. Oxford University Press, USA. ISBN 0198774508,9780198774501,0198774494,9780198774495,9780191525063.

Johansen S, Mosconi R, Nielsen B (2000). "Cointegration Analysis in the Presence of Structural Breaks in the Deterministic Trend." *Econometrics Journal*, **3**(2), 216–249. doi:10.1111/1368-423X.00047.

Johansen S, Nielsen M (2018). "The Cointegrated Vector Autoregressive Model with General Deterministic Terms." *Journal of Econometrics*, **202**(2), 214–229. doi:10.1016/j.jeconom.2017.10.003.

Juselius K (2006). *The Cointegrated VAR Model: Methodology and Applications*. Advanced Texts in Econometrics. Oxford University Press, USA. ISBN 0199285675,9780199285679.

Kilian L (1998). "Small-sample Confidence Intervals for Impulse Response Functions." *Review of Economics and Statistics*, **80**(2), 218–230. doi:10.1162/003465398557465.

Kilian L, Lütkepohl H (2017). *Structural Vector Autoregressive Analysis*. Themes in Modern Econometrics. Cambridge University Press. doi:10.1017/9781108164818.

King R, Plosser C, Stock J, Watson M (1991). "Stochastic Trends and Economic Fluctuations." *American Economic Review*, **81**(4), 819–840.

Kojadinovic I, Yan J (2010). "Modeling Multivariate Distributions with Continuous Margins using the **copula** R Package." *Journal of Statistical Software*, **34**(9), 1–20. doi:10.18637/jss.v034.i09.

Kurita T, Nielsen B (2019). "Partial Cointegrated Vector Autoregressive Models with Structural Breaks in Deterministic Terms." *Econometrics*, **7**(4), 1–35. ISSN 2225-1146. doi:10.3390/econometrics7040042.

Lange A, Dalheimer B, Herwartz H, Maxand S (2021). "**svars**: An R Package for Data-Driven Identification in Multivariate Time Series Analysis." *Journal of Statistical Software*, **97**(5), 1–34. ISSN 1548-7660. doi:10.18637/jss.v097.i05.

Larsson R, Lyhagen J, Löthgren M (2001). "Likelihood-based Cointegration Tests in Heterogeneous Panels." *Econometrics Journal*, **4**(1), 109–142. doi:10.1111/1368-423X.00059.

Lee KC, Pesaran MH (1993). "Persistence Profiles and Business Cycle Fluctuations in a Disaggregated Model of U.K. Output Growth." *Ricerche Economiche*, **47**(3), 293–322. doi:10.1016/0035-5054(93)90032-X.

Luukkonen R, Ripatti A, Saikkonen P (1999). "Testing for a Valid Normalization of Cointegrating Vectors in Vector Autoregressive Processes." *Journal of Business & Economic Statistics*, **17**(2), 195–204. doi:10.2307/1392475.

Lütkepohl H (2005). *New Introduction to Multiple Time Series Analysis*. 2nd edition. Springer. ISBN 3540401725,9783540401728.

Lütkepohl H (2006). "Structural Vector Autoregressive Analysis for Cointegrated Variables." *AStA Advances in Statistical Analysis*, **90**(1), 75–88. doi:10.1007/s10182-006-0222-4.

Lütkepohl H, Saikkonen P (2000). "Testing for the Cointegrating Rank of a VAR Process with a Time Trend." *Journal of Econometrics*, **95**(1), 177–198. doi:10.1016/S0304-4076(99)00035-4.

Lütkepohl H, Saikkonen P, Trenkler C (2004). "Testing for the Cointegrating Rank of a VAR Process with Level Shift at Unknown Time." *Econometrica*, **72**(2), 647–662. doi:10.1111/j.1468-0262.2004.00505.x.

Maddala GS, Wu S (1999). "A Comparative Study of Unit Root Tests with Panel Data and a New Simple Test." *Oxford Bulletin of Economics and Statistics*, **61**(S1), 631–652. doi:10.1111/1468-0084.0610s1631.

Matteson DS, Tsay RS (2017). "Independent Component Analysis via Distance Covariance." *Journal of the American Statistical Association*, **112**(518), 623–637. doi:10.1080/01621459.2016.1150851.

Mullen KM, Ardia D, Gil DL, Windover D, Cline J (2011). "**DEoptim**: An R Package for Global Optimization by Differential Evolution." *Journal of Statistical Software*, **40**(6), 1–26. doi:10.18637/jss.v040.i06.

Nickell S (1981). "Biases in Dynamic Models with Fixed Effects." *Econometrica*, **49**(6), 1417–26. doi:10.2307/1911408.

Onatski A (2010). "Determining the Number of Factors from Empirical Distribution of Eigenvalues." *Review of Economics and Statistics*, **92**(4), 1004–1016. doi:10.1162/REST_a_00043.

Osterwald-Lenum M (1992). "A Note with Quantiles of the Asymptotic Distribution of the Maximum Likelihood Cointegration Rank Test Statistics." *Oxford Bulletin of Economics and Statistics*, **54**(3), 461–72. doi:10.1111/j.1468-0084.1992.tb00013.x.

Pedroni P (2019). "Panel Cointegration Techniques and Open Challenges." In M Tsionas (ed.), *Panel Data Econometrics: Theory*, chapter 10, pp. 251–287. Academic Press. ISBN 978-0-12-814367-4.

Persyn D, Westerlund J (2008). "Error-Correction-Based Cointegration Tests for Panel Data." *Stata Journal*, **8**(2), 232–241. doi:10.1177/1536867X0800800205.

Pesaran MH, Shin Y (1996). "Cointegration and Speed of Convergence to Equilibrium." *Journal of Econometrics*, **71**(1-2), 117–143. doi:10.1016/0304-4076(94)01697-6.

Pesaran MH, Shin Y, Smith RJ (2000). "Structural Analysis of Vector Error Correction Models with Exogenous I(1) Variables." *Journal of Econometrics*, **97**(2), 293–343. doi:10.1016/S0304-4076(99)00073-1.

Pesaran MH, Shin Y, Smith RP (1999). "Pooled Mean Group Estimation of Dynamic Heterogeneous Panels." *Journal of the American Statistical Association*, **94**(446), 621–634. doi:10.2307/2670182.

Pesaran MH, Smith R (1995). "Estimating Long-Run Relationships from Dynamic Heterogeneous Panels." *Journal of Econometrics*, **68**(1), 79–113. doi:10.1016/0304-4076(94)01644-F.

Pfaff B (2008a). *Analysis of Integrated and Cointegrated Time Series with R*. Use R!, 2nd edition. Springer. ISBN 0387759662,9780387759661. URL https://CRAN.R-project.org/package=urca.

Pfaff B (2008b). "VAR, SVAR and SVEC Models: Implementation within R Package **vars**." *Journal of Statistical Software*, **27**(4), 1–32. doi:10.18637/jss.v027.i04.

Pretis F (2020). "Econometric Modelling of Climate Systems: The Equivalence of Energy Balance Models and Cointegrated Vector Autoregressions." *Journal of Econometrics*, **214**(1), 256–273. doi:10.1016/j.jeconom.2019.05.013.

Qu Z, Perron P (2007). "A Modified Information Criterion for Cointegration Tests Based on a VAR Approximation." *Econometric Theory*, **23**(04), 638–685. doi:10.1017/S0266466607070284.

R Core Team (2026). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Rebucci A (2010). "Estimating VARs with Long Stationary Heterogeneous Panels: A Comparison of the Fixed Effect and the Mean Group Estimators." *Economic Modelling*, **27**(5), 1183–1198. doi:10.1016/j.econmod.2010.03.001.

Risk BB, James NA, Matteson DS (2015). "**steadyICA**: ICA and Tests of Independence via Multivariate Distance Covariance." *R package version 1.0*, CRAN. URL https://CRAN.R-project.org/package=steadyICA.

Risk BB, Matteson DS, Ruppert D, Eloyan A, Caffo BS (2014). "An Evaluation of Independent Component Analyses with an Application to Resting-State fMRI." *Biometrics*, **70**(1), 224–236. ISSN 0006-341X. doi:10.1111/biom.12111.

Rizzo M, Szekely G (2022). "**energy**: E-Statistics: Multivariate Inference via the Energy of Data." *R package version 1.7-9*, CRAN. URL https://CRAN.R-project.org/package=energy.

Saikkonen P, Lütkepohl H (2000a). "Testing for the Cointegrating Rank of a VAR Process with an Intercept." *Econometric Theory*, **16**(3), 373–406. doi:10.1017/S0266466600163042.

Saikkonen P, Lütkepohl H (2000b). "Testing for the Cointegrating Rank of a VAR Process with Structural Shifts." *Journal of Business & Economic Statistics*, **18**(4), 451–64. doi:10.1080/07350015.2000.10524884.

Saikkonen P, Lütkepohl H (2000c). "Trend Adjustment Prior to Testing for the Cointegrating Rank of a Vector Autoregressive Process." *Journal of Time Series Analysis*, **21**(4), 435–456. doi:10.1111/1467-9892.00192.

Sanchez-Martinez M, Woźniak T (2026). "Forecasting with Bayesian Panel Vector Autoregressions: Using the R Package **bpvars**." arXiv:2606.14143 [econ.EM]. doi:10.48550/arXiv.2606.14143.

Sharpsteen C, Bracken C (2020). "**tikzDevice**: R Graphics Output in LaTeX Format." *R package version 0.12.3.1*, CRAN. URL https://CRAN.R-project.org/package=tikzDevice.

Sigmund M, Ferstl R (2019). "Panel Vector Autoregression in R with the Package **Panelvar**." *Quarterly Review of Economics and Finance*. doi:10.2139/ssrn.2896087.

Sims CA (1980). "Macroeconomics and Reality." *Econometrica*, **48**(1), 1–48. doi:10.2307/1912017.

Smyth R, Narayan P (2015). "Applied Econometrics and Implications for Energy Economics Research." *Energy Economics*, **50**(C), 351–358. doi:10.1016/j.eneco.2014.07.023.

StataCorp (2025). *Stata Statistical Software: Release 19*. StataCorp LLC, College Station, Texas. URL https://www.stata.com/.

Stouffer SA, Suchman EA, DeVinney LC, Star SA, Williams RM (1949). *The American Soldier*, volume 1 of *Studies in Social Psychology in World War II*. Princeton University Press. doi:10.1177/000271624926500124.

Székely GJ, Rizzo ML, Bakirov NK (2007). "Measuring and Testing Dependence by Correlation of Distances." *Annals of Statistics*, **35**(6), 2769–2794. doi:10.1214/009053607000000505.

Tang Y, Horikoshi M, Wenxuan L (2016). "**ggfortify**: Unified Interface to Visualize Statistical Results of Popular R Packages." *The R Journal*, **8**(2), 474–485. doi:10.32614/RJ-2016-060.

Trenkler C (2008). "Determining p-Values for Systems Cointegration Tests with a Prior Adjustment for Deterministic Terms." *Computational Statistics*, **23**(1), 19–39. doi:10.1007/s00180-007-0066-8.

Trenkler C, Saikkonen P, Lütkepohl H (2008). "Testing for the Cointegrating Rank of a VAR Process with Level Shift and Trend Break." *Journal of Time Series Analysis*, **29**(2), 331–358. doi:10.1111/j.1467-9892.2007.00558.x.

Venables WN, Ripley BD (2002). *Modern Applied Statistics with S*. 4th edition. Springer. ISBN 0-387-95457-0. URL https://www.stats.ox.ac.uk/pub/MASS4/.

Vlaar PJ (2004). "On the Asymptotic Distribution of Impulse Response Functions with Long-Run Restrictions." *Econometric Theory*, **20**(5), 891–903. doi:10.1017/S0266466604205047.

Wickham H (2011). "**testthat**: Get Started with Testing." *The R Journal*, **3**(1), 5–10. doi:10.32614/RJ-2011-002.

Wickham H (2016). ***ggplot2**: Elegant Graphics for Data Analysis*. Use R!, 2nd edition. Springer. ISBN 331924275X,9783319242750. URL https://ggplot2.tidyverse.org.

Yang M (2002). "Lag Length and Mean Break in Stationary VAR Models." *Econometrics Journal*, **5**(2), 374–387. doi:10.1111/1368-423X.00089.

Örsal DDK (2017). "Analysing Money Demand Relation for OECD Countries Using Common Factors." *Applied Economics*, **49**(60), 6003–6013. doi:10.1080/00036846.2017.1371842.

Örsal DDK, Arsova A (2017). "Meta-Analytic Cointegrating Rank Tests for Dependent Panels." *Econometrics and Statistics*, **2**(C), 61–72. doi:10.1016/j.ecosta.2016.10.001.

Örsal DDK, Droge B (2011). "Corrigendum to 'Likelihood-Based Cointegration Tests in Heterogeneous Panels' (Larsson R., J. Lyhagen and M. Löthgren, Econometrics Journal, 4, 2001, 109–142)." *Econometrics Journal*, **14**(1), 121–125. doi:10.1111/j.1368-423X.2010.00327.x.

Örsal DDK, Droge B (2014). "Panel Cointegration Testing in the Presence of a Time Trend." *Computational Statistics & Data Analysis*, **76**(C), 377–390. doi:10.1016/j.csda.2012.05.017.

Affiliation:

Lennart Empting
Faculty of Economics
Universität Duisburg-Essen
Universitätsstraße 12
45141 Essen, Germany
E-mail: lennart.empting@vwl.uni-due.de
URL: https://github.com/Lenni89